



Cancer By Site

Percent Distribution of Cancer

2006 Total Cancer Incidence of Washington State Residents




2006 Total Cancer Mortality of Washington State Residents




Percent Distribution of Cancer

2006 Male Cancer Incidence of Washington State Residents




2006 Male Cancer Mortality of Washington State Residents




Percent Distribution of Cancer

2006 Female Cancer Incidence of Washington State Residents



2006 Female Cancer Mortality of Washington State Residents



All Sites Combined

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using ICD-03 codes C00.0 - C80.9, all histology codes. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 codes 140-208 (1992-1998), ICD-10 codes C00-C97 (1999-2006).

2006 Washington State Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


Incidence			
	Male	Female	Total
All Reportable Cancers	16,975	16,833	33,810
2006 Incidence Rates	591.9	492.7	531.6
2005 National Rates	551.2	456.6	492.8

Death			
	Male	Female	Total
Deaths	5,606	5,397	11,003
2006 Death Rates	208.6	154.1	176.1
2005 National Rates	226.1	155.3	184.0

Stage at Diagnosis

2006 Washington State Residents

Total Incidence and Percents



Relative Survival Rates

2005 National 5-year rates¹

In situ	n/a
Local	n/a
Regional	n/a
Distant	n/a
Unstaged	n/a
Overall	68%


n/a = not available

Figures include in situ and invasive cancers

¹ 5-Year Relative Survival Rates for cases diagnosed 2000-2004 based on follow-up of patients through 2005. Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.


All Sites Combined

Washington State Residents by Gender
Incidence Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined, cancer incidence rates for all cancer sites combined increased slightly by 0.7% per year from 1992 to 2001; since then, however the rates have remained level.
- For men, cancer incidence rates for all cancer sites combined decreased steadily by 3.4% per year from 1992 to 1995; rates remained level from 1995 to 1999; since 1999 the rates have decreased slightly by 0.8% per year.
- For women, cancer incidence rates for all cancer sites combined increased steadily by 2.1% per year from 1992 and 1998; since then, however the rates have remained level.

Washington State Residents by Gender
Death Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined, cancer mortality rates for all cancer sites combined decreased slightly by 0.9% per year from 1992 to 2003; since 2003 the rates have decreased steadily by 2.4% per year.
- For men, cancer mortality rates for all cancer sites combined decreased steadily by 1.5% per year during the period from 1992 to 2006.
- For women, cancer mortality rates for all cancer sites combined decreased slightly by 0.7% per year from 1992 to 2003; since 2003 the rates have decreased steadily by 2.2% per year.


The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.
 Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.

All Sites Combined

2004-2006 Washington State Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population




NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.
Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2006 released 1/2009.

All Sites Combined

2004-2006 Washington State Residents

Average Annual Death Rates by County with rates in [] and 95% CI's |—|

Age-adjusted to 2000 US standard population



NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.

Bladder

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using primary site ICD-03 codes C67.0-C67.9, excluding histology codes 9140, 9590-9989. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 codes 188.0-188.9 (1992-1998), ICD-10 code C67 (1999-2006).

2006 Washington State Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


Incidence			
	Male	Female	Total
All Reportable Cancers	1,095	326	1,422
2006 Incidence Rates	40.7	9.4	23.0
2005 National Rates	36.8	9.2	20.9

Death			
	Male	Female	Total
Deaths	203	85	288
2006 Death Rates	8.1	2.3	4.6
2005 National Rates	7.6	2.2	4.3

Stage at Diagnosis

2006 Washington State Residents

Total Incidence and Percents



Relative Survival Rates

2005 National 5-year rates¹

In situ	n/a
Local	93%
Regional	44%
Distant	6%
Unstaged	56%
Overall	80%


n/a = not available

Figures include in situ and invasive cancers

¹ 5-Year Relative Survival Rates for cases diagnosed 2000-2004 based on follow-up of patients through 2005. Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.


Bladder

Washington State Residents by Gender
Incidence Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined and women separately, bladder cancer incidence rates remained level during the period from 1992 to 2006.
- For men, bladder cancer incidence rates increased steadily by 2.2% per year from 1992 to 1998; since then, however the rates have remained level.

Washington State Residents by Gender
Death Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined and men and women separately, bladder cancer mortality rates remained level during the period from 1992 to 2006.


The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.
 Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.

Bladder

2004-2006 Washington State Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population




NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.
Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2006 released 1/2009.

Bladder

2004-2006 Washington State Residents

Average Annual Death Rates by County with rates in [] and 95% CI's |—|

Age-adjusted to 2000 US standard population



NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.

Brain

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using primary site ICD-03 codes C71.0-C71.9, excluding histology codes 9530-9539 and 9590-9989. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 codes 191.0-191.9 (1992-1998), ICD-10 code C71 (1999-2006).

2006 Washington State Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


Incidence			
	Male	Female	Total
All Reportable Cancers	231	177	408
2006 Incidence Rates	7.5	5.3	6.3
2005 National Rates	7.1	5.0	6.0

Death			
	Male	Female	Total
Deaths	202	122	324
2006 Death Rates	6.8	3.5	5.0
2005 National Rates	5.3	3.5	4.3

Stage at Diagnosis

2006 Washington State Residents

Total Incidence and Percents



Relative Survival Rates

2005 National 5-year rates¹

In situ	n/a
Local	n/a
Regional	n/a
Distant	n/a
Unstaged	n/a
Overall	31%

n/a - not available

Figures include only invasive cancers


¹ 5-Year Relative Survival Rates for cases diagnosed 2000-2004 based on follow-up of patients through 2005. Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.

Brain

Washington State Residents by Gender

Incidence Rate Trends

Age-adjusted to 2000 US standard population




- For both sexes combined and men and women separately, brain cancer incidence rates remained level during the period from 1992 to 2006.

Washington State Residents by Gender

Death Rate Trends

Age-adjusted to 2000 US standard population



- For both sexes combined and men and women separately, brain cancer mortality rates remained level during the period from 1992 to 2006.

The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.


Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.

Brain

2004-2006 Washington State Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population




NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.
Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2006 released 1/2009.

Brain

2004-2006 Washington State Residents

Average Annual Death Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population



NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.

Breast (Female)

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using primary site ICD-03 codes C50.0-C50.9, excluding histology codes 9140, 9590-9989. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 codes 174.0-174.9 (1992-1998), ICD-10 codes C50 (1999-2006).

2006 Washington State Female Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


Incidence			
	Male	Female	Total
All Reportable Cancers	n/a	5,531	n/a
2006 Incidence Rates	n/a	160.1	n/a
2005 National Rates	n/a	151.1	n/a

Death			
	Male	Female	Total
Deaths	n/a	809	n/a
2006 Death Rates	n/a	22.9	n/a
2005 National Rates	n/a	24.0	n/a

Stage at Diagnosis

2006 Washington State Female Residents

Total Incidence and Percents



Relative Survival Rates	
2005 National 5-year rates ¹	
In situ	100%
Local	98%
Regional	84%
Distant	28%
Unstaged	56%
Overall	91%

Figures include in situ and invasive cancers

¹ 5-Year Relative Survival Rates for cases diagnosed 2000-2004 based on follow-up of patients through 2005.

n/a Not available for sex-specific sites.


Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.

Breast (Female)

Washington State Female Residents

Incidence Rate Trends

Age-adjusted to 2000 US standard population




- For women, breast cancer incidence rates increased steadily by 2.9% per year from 1992 to 1999; since then, however the rates have decreased steadily by 2.1% per year.

Washington State Female Residents

Death Rate Trends

Age-adjusted to 2000 US standard population



- For women, breast cancer mortality rates decreased steadily by 3.3% per year from 1992 to 2000; since then, however the rates have remained level.

The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.


Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.

Breast (Female)

2004-2006 Washington State Female Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population




NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.
Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2006 released 1/2009.

Breast (Female)

2004-2006 Washington State Female Residents

Average Annual Death Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population



NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.

Cervix (Uterine)

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using primary site ICD-03 codes C53.0-C53.9, excluding histology codes 9140, 9590-9989 and excluding in situ. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 codes 180.0-180.9 (1992-1998), ICD-10 code C53 (1999-2006).

2006 Washington State Female Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


Incidence			
	Male	Female	Total
All Reportable Cancers	n/a	244	n/a
2006 Incidence Rates	n/a	7.6	n/a
2005 National Rates	n/a	8.0	n/a

Death			
	Male	Female	Total
Deaths	n/a	69	n/a
2006 Death Rates	n/a	2.0	n/a
2005 National Rates	n/a	2.4	n/a

Stage at Diagnosis

2006 Washington State Female Residents

Total Incidence and Percents



Relative Survival Rates

2005 National 5-year rates¹

In situ	n/a
Local	91%
Regional	57%
Distant	17%
Unstaged	53%
Overall	71%

n/a - not available

Figures include only invasive cancers


¹ 5-Year Relative Survival Rates for cases diagnosed 2000-2004 based on follow-up of patients through 2005.

n/a Not available for sex-specific sites.

Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.


Cervix (Uterine)

Washington State Female Residents
Incidence Rate Trends
 Age-adjusted to 2000 US standard population



- For women, cervical cancer incidence rates decreased steadily by 2.2% per year during the period from 1992 to 2006.

Washington State Female Residents
Death Rate Trends
 Age-adjusted to 2000 US standard population



- For women, cervical cancer mortality rates decreased steadily by 2.6% per year during the period from 1992 to 2006.


The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.
 Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.

Cervix (Uterine)

2004-2006 Washington State Female Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population




NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.
Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2006 released 1/2009.

Cervix (Uterine)

2004-2006 Washington State Female Residents

Average Annual Death Rates by County with rates in [] and 95% CI's |—|

Age-adjusted to 2000 US standard population



NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.

Colorectal

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using primary site ICD-03 codes C18.0-C18.9, C19.9, C20.9, C26.0, excluding histology codes 9140, 9590-9989. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 codes 153.0-154.1, 159.0 (1992-1998), ICD-10 codes C18-C20, C26.0 (1999-2006).

2006 Washington State Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


Incidence			
	Male	Female	Total
All Reportable Cancers	1,416	1,321	2,737
2006 Incidence Rates	50.4	37.9	43.5
2005 National Rates	58.4	43.2	50.0

Death			
	Male	Female	Total
Deaths	473	451	924
2006 Death Rates	17.4	12.5	14.7
2005 National Rates	21.0	14.6	17.4

Stage at Diagnosis

2006 Washington State Residents

Total Incidence and Percents



Relative Survival Rates

2005 National 5-year rates¹

In situ	96%
Local	90%
Regional	69%
Distant	11%
Unstaged	37%
Overall	67%

Figures include in situ and invasive cancers


¹ 5-Year Relative Survival Rates for cases diagnosed 2000-2004 based on follow-up of patients through 2005. Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.

Colorectal

Washington State Residents by Gender

Incidence Rate Trends

Age-adjusted to 2000 US standard population




- For both sexes combined, colorectal cancer incidence rates remained level from 1992 to 1998; since then, however the rates have decreased steadily by 3.3% per year.
- For men, colorectal cancer incidence rates remained level from 1992 to 1998; since then, however the rates have decreased steadily by 3.7% per year.
- For women, colorectal cancer incidence rates remained level from 1992 to 1998; since then, however the rates have decreased steadily by 3.1% per year.

Washington State Residents by Gender

Death Rate Trends

Age-adjusted to 2000 US standard population



- For both sexes combined and women separately, colorectal cancer mortality rates decreased steadily by 2.6% per year during the period from 1992 to 2006.
- For men, colorectal cancer mortality rates decreased steadily by 2.7% per year during the period from 1992 to 2006.

The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.


Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.

Colorectal

2004-2006 Washington State Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population




NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.
Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2006 released 1/2009.

Colorectal

2004-2006 Washington State Residents

Average Annual Death Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population



NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.

Endometrium

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using primary site ICD-03 codes C54.0-C54.9, excluding histology codes 9140, 9590-9989. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 codes 182.0-182.1, 182.8 (1992-1998), ICD-10 codes C54 (1999-2006).

2006 Washington State Female Residents Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


Incidence			
	Male	Female	Total
All Reportable Cancers	n/a	812	n/a
2006 Incidence Rates	n/a	23.1	n/a
2005 National Rates	n/a	23.4	n/a

Death			
	Male	Female	Total
Deaths	n/a	73	n/a
2006 Death Rates	n/a	2.2	n/a
2005 National Rates	n/a	1.9	n/a

Stage at Diagnosis

2006 Washington State Female Residents

Total Incidence and Percents



Relative Survival Rates	
2005 National 5-year rates ¹	
In situ	99%
Local	96%
Regional	68%
Distant	25%
Unstaged	58%
Overall	84%

n/c not calculated because the number of cases is less than 5

Figures include in situ and invasive cancers


¹ 5-Year Relative Survival Rates for cases diagnosed 2000-2004 based on follow-up of patients through 2005.

n/a Not available for sex-specific sites.

Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.


Endometrium

Washington State Female Residents
Incidence Rate Trends
 Age-adjusted to 2000 US standard population



- For women, endometrial cancer incidence rates decreased steadily by 1.0% per year during the period from 1992 to 2006.

Washington State Female Residents
Death Rate Trends
 Age-adjusted to 2000 US standard population



- For women, endometrial cancer mortality rates remained level during the period from 1992 to 2006.

The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.


Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.

Endometrium

2004-2006 Washington State Female Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population




NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.
Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2006 released 1/2009.

Endometrium

2004-2006 Washington State Female Residents

Average Annual Death Rates by County with rates in [] and 95% CI's |—|

Age-adjusted to 2000 US standard population



NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.

Esophagus

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using primary site ICD-03 codes C15.0-C15.9, excluding histology codes 9140, 9590-9989. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 codes 150.0-150.9 (1992-1998), ICD-10 code C15 (1999-2006).

2006 Washington State Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


Incidence			
	Male	Female	Total
All Reportable Cancers	231	82	313
2006 Incidence Rates	8.0	2.4	5.0
2005 National Rates	7.7	1.8	4.4

Death			
	Male	Female	Total
Deaths	234	81	315
2006 Death Rates	8.4	2.3	5.0
2005 National Rates	7.9	1.7	4.4

Stage at Diagnosis

2006 Washington State Residents

Total Incidence and Percents



Relative Survival Rates

2005 National 5-year rates¹

In situ	72%
Local	35%
Regional	18%
Distant	3%
Unstaged	12%
Overall	17%


n/c - not calculated because the number of cases is less than 5

Figures include in situ and invasive cancers

¹ 5-Year Relative Survival Rates for cases diagnosed 2000-2004 based on follow-up of patients through 2005. Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.


Esophagus

Washington State Residents by Gender
Incidence Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined, esophageal cancer incidence rates increased steadily by 3.1% per year from 1992 to 2000; since then, however the rates have decreased steadily by 3.5% per year.
- For men, esophageal cancer incidence rates increased sharply by 4.1% per year from 1992 to 1999, since then, however the rates have decreased steadily by 3.5% per year.
- For women, esophageal cancer incidence rates remained level during the period from 1992 to 2006.

Washington State Residents by Gender
Death Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined, esophageal cancer mortality rates remained level during the period from 1992 to 2006.
- For men, esophageal cancer mortality rates increased steadily by 1.0% per year during the period from 1992 to 2006.
- For women, esophageal cancer mortality rates remained level from 1992 to 2004; since then, however the rates have increased sharply by 21.5% per year.

The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.


Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.

Esophagus

2004-2006 Washington State Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population




NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.
 Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2006 released 1/2009.

Esophagus

2004-2006 Washington State Residents

Average Annual Death Rates by County with rates in [] and 95% CI's |—|

Age-adjusted to 2000 US standard population



NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.

Hodgkin Lymphoma

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using all primary sites with histology codes 9650-9667. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 codes 201.0-201.9 (1992-1998), ICD-10 code C81 (1999-2006).

2006 Washington State Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


Incidence			
	Male	Female	Total
All Reportable Cancers	97	77	174
2006 Incidence Rates	3.1	2.4	2.7
2005 National Rates	3.2	2.6	2.8

Death			
	Male	Female	Total
Deaths	14	9	23
2006 Death Rates	0.5	0.3	0.4
2005 National Rates	0.5	0.4	0.4

Stage at Diagnosis

2006 Washington State Residents

Total Incidence and Percents



Relative Survival Rates

2005 National 5-year rates¹

In situ	n/a
Local	n/a
Regional	n/a
Distant	n/a
Unstaged	n/a
Overall	85%

n/a - not available


n/c - not calculated because the number of cases is less than 5

Figures include only invasive cancers

¹ 5-Year Relative Survival Rates for cases diagnosed 2000-2004 based on follow-up of patients through 2005. Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.


Hodgkin Lymphoma

Washington State Residents by Gender
Incidence Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined and men and women separately, incidence rates for Hodgkin Lymphoma remained level during the period from 1992 to 2006.

Washington State Residents by Gender
Death Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined and men and women separately, mortality rates for Hodgkin Lymphoma remained level during the period from 1992 to 2006.


The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.
 Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.

Hodgkin Lymphoma

2004-2006 Washington State Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |—|

Age-adjusted to 2000 US standard population




NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.
Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2006 released 1/2009.

Hodgkin Lymphoma

2004-2006 Washington State Residents

Average Annual Death Rates by County with rates in [] and 95% CI's |—|

Age-adjusted to 2000 US standard population



NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.

Kaposi Sarcoma

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using all primary sites with histology code 9140. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 code 176 (1992-1998), ICD-10 codes C46 (1999-2006).

2006 Washington State Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


Incidence			
	Male	Female	Total
All Reportable Cancers	27	3	30
2006 Incidence Rates	0.9	n/c	0.5
2005 National Rates	1.2	0.1	0.6

Death			
	Male	Female	Total
Deaths	0	0	0
2006 Death Rates	0.0	0.0	0.0
2005 National Rates	0.0	0.0	0.0

Stage at Diagnosis

2006 Washington State Residents

Total Incidence and Percents



Relative Survival Rates	
2005 National 5-year rates ¹	
In situ	n/a
Local	n/a
Regional	n/a
Distant	n/a
Unstaged	n/a
Overall	61%


n/a - not available

n/c - not calculated because the number of cases is less than 5
 Figures include only invasive cancers

¹ 5-Year Relative Survival Rates for cases diagnosed 2000-2004 based on follow-up of patients through 2005. Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.


Kaposi Sarcoma

Washington State Residents by Gender
Incidence Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined and men separately, incidence rates for Kaposi's Sarcoma remained level from 1992 to 1995; rates decreased sharply by 42.3% per year from 1995 to 1998; since 1998 the rates have remained level.
- Trends in incidence rates for Kaposi's Sarcoma from 1992 to 2006 could not be assessed for women due to small numbers of new cases from Kaposi's Sarcoma in this group.

Washington State Residents by Gender
Death Rate Trends
 Age-adjusted to 2000 US standard population



- Trends in mortality rates for Kaposi's Sarcoma from 1992 to 2006 could not be assessed for men and women combined or separately due to small numbers of deaths from Kaposi's Sarcoma in these groups.

The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.

Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.

Kaposi Sarcoma

2004-2006 Washington State Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population



NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.
Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2006 released 1/2009.

Kaposi Sarcoma

2004-2006 Washington State Residents

Average Annual Death Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population



NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.

Kidney and Renal Pelvis

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using primary site ICD-03 codes C64.9, C65.9, excluding histology codes 9140, 9590-9989. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 codes 189.0, 189.1 (1992-1998), ICD-10 codes C64-C65 (1999-2006).

2006 Washington State Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


Incidence			
	Male	Female	Total
All Reportable Cancers	604	389	993
2006 Incidence Rates	19.9	11.5	15.4
2005 National Rates	19.3	9.6	14.0

Death			
	Male	Female	Total
Deaths	131	97	228
2006 Death Rates	4.6	2.7	3.5
2005 National Rates	5.9	2.7	4.1

Stage at Diagnosis

2006 Washington State Residents

Total Incidence and Percents




Relative Survival Rates	
2005 National 5-year rates ¹	
In situ	87%
Local	90%
Regional	61%
Distant	10%
Unstaged	37%
Overall	68%

Figures include in situ and invasive cancers

¹ 5-Year Relative Survival Rates for cases diagnosed 2000-2004 based on follow-up of patients through 2005. Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.


Kidney and Renal Pelvis

Washington State Residents by Gender
Incidence Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined, kidney and renal pelvis cancer increased steadily by 2.5% per year during the period from 1992 to 2006.
- For men, kidney and renal pelvis cancer increased steadily by 2.3% per year during the period from 1992 to 2006 .
- For women, kidney and renal pelvis cancer remained level from 1992 to 1994; since then, however the rates have increased steadily by 3.5% per year.

Washington State Residents by Gender
Death Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined and men and women separately, kidney and renal pelvis cancer mortality rates remained level during the period from 1992 to 2006.

The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.


Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.

Kidney and Renal Pelvis

2004-2006 Washington State Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population




NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.
Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2006 released 1/2009.

Kidney and Renal Pelvis

2004-2006 Washington State Residents

Average Annual Death Rates by County with rates in [] and 95% CI's |—|

Age-adjusted to 2000 US standard population



NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.

Larynx

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using primary site ICD-03 codes C32.0-C32.9, excluding histology codes 9140, 9590-9989. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 codes 161.0-161.9 (1992-1998), ICD-10 code C32 (1999-2006).

2006 Washington State Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


Incidence			
	Male	Female	Total
All Reportable Cancers	154	50	204
2006 Incidence Rates	5.3	1.5	3.2
2005 National Rates	6.6	1.3	3.7

Death			
	Male	Female	Total
Deaths	36	18	54
2006 Death Rates	1.3	0.5	0.9
2005 National Rates	2.2	0.5	1.2

Stage at Diagnosis

2006 Washington State Residents

Total Incidence and Percents



Relative Survival Rates

2005 National 5-year rates¹


In situ	95%
Local	81%
Regional	50%
Distant	21%
Unstaged	44%
Overall	64%

Figures include in situ and invasive cancers

¹ 5-Year Relative Survival Rates for cases diagnosed 2000-2004 based on follow-up of patients through 2005. Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.


Larynx

Washington State Residents by Gender Incidence Rate Trends Age-adjusted to 2000 US standard population



- For both sexes combined, laryngeal cancer incidence rates decreased steadily by 2.8% per year during the period from 1992 to 2006.
- For men, laryngeal cancer incidence rates decreased steadily by 3.0% per year during the period from 1992 to 2006.
- For women, laryngeal cancer incidence rates decreased steadily by 1.9% per year during the period from 1992 to 2006.

Washington State Residents by Gender Death Rate Trends Age-adjusted to 2000 US standard population



- For both sexes combined, laryngeal cancer mortality rates decreased steadily by 1.6% per year during the period from 1992 to 2006.
- For men, laryngeal cancer mortality rates decreased steadily by 2.1% per year during the period from 1992 to 2006.
- For women, laryngeal cancer mortality rates remained level during the period from 1992 to 2006.

The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.


Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.

Larynx

2004-2006 Washington State Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population




NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.
Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2006 released 1/2009.

Larynx

2004-2006 Washington State Residents

Average Annual Death Rates by County with rates in [] and 95% CI's |—|

Age-adjusted to 2000 US standard population



NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.

Leukemia

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using all sites with histology 9733, 9742, 9800-9820, 9826, 9831-9948, 9963,9964; for sites C420, C421 and C424 only, histology 9823 or 9827. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 codes 204.0-208.9 (1992-1998), ICD-10 codes C90.1, C91-C95 (1999-2006).

2006 Washington State Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


Incidence			
	Male	Female	Total
All Reportable Cancers	489	379	868
2006 Incidence Rates	17.2	11.2	13.8
2005 National Rates	14.8	8.8	11.4

Death			
	Male	Female	Total
Deaths	243	194	437
2006 Death Rates	9.3	5.4	7.1
2005 National Rates	9.7	5.4	7.2

Stage at Diagnosis

2006 Washington State Residents

Total Incidence and Percents



Relative Survival Rates	
2005 National 5-year rates ¹	
In situ	n/a
Local	n/a
Regional	n/a
Distant	52%
Unstaged	n/a
Overall	52%


n/a - not available

n/c - not calculated because the number of cases is less than 5
 Figures include only invasive cancers

¹ 5-Year Relative Survival Rates for cases diagnosed 2000-2004 based on follow-up of patients through 2005. Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.


Leukemia

Washington State Residents by Gender
Incidence Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined, leukemia incidence rates remained level from 1992 to 1994; since then, however the rates decreased slightly by 0.6% per year.
- For men and women separately, leukemia incidence rates remained level during the period from 1992 to 2006.

Washington State Residents by Gender
Death Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined, leukemia mortality rates decreased slightly by 0.8% per year during the period from 1992 to 2006.
- For men and women separately, leukemia mortality rates remained level during the period from 1992 to 2006.


The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.
 Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.

Leukemia

2004-2006 Washington State Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population




NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.
Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2006 released 1/2009.

Leukemia

2004-2006 Washington State Residents

Average Annual Death Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population



NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.

Liver

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using primary site ICD-03 code C22.0, excluding histology codes 9140, 9590-9989. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 codes 155.0, 155.2 (1992-1998), ICD-10 codes C22.0, C22.2-C22.4, C22.7, C22.9 (1999-2006).

2006 Washington State Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


Incidence			
	Male	Female	Total
All Reportable Cancers	273	104	377
2006 Incidence Rates	8.7	3.1	5.8
2005 National Rates	9.7	3.2	6.2

Death			
	Male	Female	Total
Deaths	188	70	258
2006 Death Rates	6.2	2.0	4.0
2005 National Rates	6.4	2.2	4.1

Stage at Diagnosis

2006 Washington State Residents

Total Incidence and Percents



Relative Survival Rates

2005 National 5-year rates¹

In situ	n/a
Local	26%
Regional	8%
Distant	3%
Unstaged	6%
Overall	13%


n/a - not available

Figures include only invasive cancers

¹ 5-Year Relative Survival Rates for cases diagnosed 2000-2004 based on follow-up of patients through 2005. Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.


Liver

Washington State Residents by Gender
Incidence Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined, liver cancer incidence rates increased sharply by 5.0% per year during the period from 1992 to 2006.
- For men, liver cancer incidence rates increased sharply by 4.9% per year during the period from 1992 to 2006.
- For women, liver cancer incidence rates increased sharply by 4.1% per year during the period from 1992 to 2006.

Washington State Residents by Gender
Death Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined, liver cancer mortality rates increased steadily by 3.4% per year during the period from 1992 to 2006.
- For men, liver cancer mortality rates increased steadily by 3.5% per year during the period from 1992 to 2006.
- For women, liver cancer mortality rates increased steadily by 2.5% per year during the period from 1992 to 2006.

The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.


Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.

Liver

2004-2006 Washington State Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population




NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.
 Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2006 released 1/2009.

Liver

2004-2006 Washington State Residents

Average Annual Death Rates by County with rates in [] and 95% CI's |—|

Age-adjusted to 2000 US standard population



NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.

Lung and Bronchus

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using primary site ICD-03 codes C34.0-C34.9, excluding histology codes 9140, 9590-9989. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 codes 162.2-162.9 (1992-1998), ICD-10 code C34 (1999-2001).

2006 Washington State Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


Incidence			
	Male	Female	Total
All Reportable Cancers	2,067	1,996	4,063
2006 Incidence Rates	75.5	60.1	66.4
2005 National Rates	74.1	51.7	61.1

Death			
	Male	Female	Total
Deaths	1,631	1,441	3,072
2006 Death Rates	60.8	42.4	50.2
2005 National Rates	69.4	40.6	52.8

Stage at Diagnosis

2006 Washington State Residents

Total Incidence and Percents




Relative Survival Rates	
2005 National 5-year rates ¹	
In situ	44%
Local	50%
Regional	21%
Distant	3%
Unstaged	8%
Overall	15%

n/c - not calculated because the number of cases is less than 5
 Figures include in situ and invasive cancers

¹ 5-Year Relative Survival Rates for cases diagnosed 2000-2004 based on follow-up of patients through 2005. Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.


Lung and Bronchus

Washington State Residents by Gender
Incidence Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined, lung and bronchus cancer incidence rates remained level from 1992 to 1998; since then, however the rates have decreased steadily by 1.3% per year.
- For men, lung and bronchus cancer incidence rates decreased steadily by 1.6% per year during the period from 1992 to 2006.
- For women, lung and bronchus cancer incidence rates increased sharply by 4.9% per year from 1992 to 1995; since then, however the rates have remained level.

Washington State Residents by Gender
Death Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined, lung and bronchus cancer mortality rates decreased steadily by 1.2% per year during the period from 1992 to 2006.
- For men, lung and bronchus cancer mortality rates decreased steadily by 2.2% per year during the period from 1992 to 2006.
- For women, lung and bronchus cancer mortality rates remained level from 1992 to 2003; since then, however the rates have decreased steadily by 3.5% per year.


The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.
 Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.

Lung and Bronchus

2004-2006 Washington State Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population




NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.
Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2006 released 1/2009.

Lung and Bronchus

2004-2006 Washington State Residents

Average Annual Death Rates by County with rates in [] and 95% CI's |—|

Age-adjusted to 2000 US standard population



NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.

Melanoma of the Skin

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using primary site ICD-03 codes C44.0-C44.9, including only histology codes 8720-8790. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 codes 172.0-172.9 (1992-1998), ICD-10 code C43 (1999-2006).

2006 Washington State Residents

Incidence and Death Summary by Sex


Counts and rates per 100,000 adjusted to the 2000 standard population

Incidence			
	Male	Female	Total
All Reportable Cancers	1,473	1,290	2,763
2006 Incidence Rates	49.8	38.8	43.0
2005 National Rates	43.3	27.6	34.0

Death			
	Male	Female	Total
Deaths	115	75	190
2006 Death Rates	4.0	2.1	3.0
2005 National Rates	4.0	1.8	2.7

Stage at Diagnosis

2006 Washington State Residents




Relative Survival Rates	
2005 National 5-year rates ¹	
In situ	100%
Local	99%
Regional	65%
Distant	15%
Unstaged	76%
Overall	96%

Figures include in situ and invasive cancers

¹ 5-Year Relative Survival Rates for cases diagnosed 2000-2004 based on follow-up of patients through 2005. Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.


Melanoma of the Skin

Washington State Residents by Gender
Incidence Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined, melanoma incidence rates increased sharply by 11.9% per year from 1992 to 1995; since 1995 the rates have increased steadily by 3.5% per year.
- For men, melanoma incidence rates increased sharply by 4.3% per year during the period from 1992 to 2006.
- For women, melanoma incidence rates increased sharply by 13.1% per year from 1992 to 1995; since 1995 the rates have increased steadily by 3.5% per year.

Washington State Residents by Gender
Death Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined and men and women separately, melanoma mortality rates remained level during the period from 1992 to 2006.

The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.


Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.

Melanoma of the Skin

2004-2006 Washington State Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population




NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.
 Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2006 released 1/2009.

Melanoma of the Skin

2004-2006 Washington State Residents

Average Annual Death Rates by County with rates in [] and 95% CI's |—|

Age-adjusted to 2000 US standard population



NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.

Multiple Myeloma

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using all primary sites with histology codes 9731-9732, 9734. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 codes 203.0, 203.2-203.8 (1992-1998), ICD-10 codes C90.0, C90.2 (1999-2006).

2006 Washington State Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


Incidence			
	Male	Female	Total
All Reportable Cancers	212	154	366
2006 Incidence Rates	7.6	4.5	5.8
2005 National Rates	6.9	4.6	5.6

Death			
	Male	Female	Total
Deaths	115	89	204
2006 Death Rates	4.4	2.5	3.3
2005 National Rates	4.5	2.9	3.6

Stage at Diagnosis

2006 Washington State Residents

Total Incidence and Percents



Relative Survival Rates

2005 National 5-year rates¹

In situ	n/a
Local	n/a
Regional	n/a
Distant	36%
Unstaged	n/a
Overall	36%

n/a - not available


n/c - not calculated because the number of cases is less than 5

Figures include only invasive cancers

¹ 5-Year Relative Survival Rates for cases diagnosed 2000-2004 based on follow-up of patients through 2005. Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.


Multiple Myeloma

Washington State Residents by Gender
Incidence Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined, multiple myeloma incidence rates remained level from 1992 to 1996; since then, however the rates have decreased steadily by 1.3% per year.
- For men and women separately, multiple myeloma incidence rates remained level during the period from 1992 to 2006.

Washington State Residents by Gender
Death Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined, multiple myeloma mortality rates remained level from 1992 to 2002; since then, however the rates have decreased sharply by 6.9% per year.
- For men, multiple myeloma mortality rates remained level from 1992 to 2002; since then, however the rates have decreased sharply by 5.4% per year.
- For women, multiple myeloma mortality rates remained level during the period from 1992 to 2006.

The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.


Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.

Multiple Myeloma

2004-2006 Washington State Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population




NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.
 Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2006 released 1/2009.

Multiple Myeloma

2004-2006 Washington State Residents

Average Annual Death Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population



NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.

Non-Hodgkin Lymphoma

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry all sites with histology 9590-9596, 9670-9729; and except for sites C420, C421 and C424 all other sites with histology 9823 or 9827. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 codes 200.0-200.8, 202.0-202.2, 202.8-202.9 (1992-1998), ICD-10 codes C82-C85 (1999-2006).

2006 Washington State Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


Incidence			
	Male	Female	Total
All Reportable Cancers	780	596	1,376
2006 Incidence Rates	27.4	17.4	21.8
2005 National Rates	23.4	16.0	19.2

Death			
	Male	Female	Total
Deaths	215	212	427
2006 Death Rates	8.1	5.9	6.8
2005 National Rates	8.8	5.5	6.9

Stage at Diagnosis

2006 Washington State Residents

Total Incidence and Percents



Relative Survival Rates

2005 National 5-year rates¹

In situ	n/a
Local	n/a
Regional	n/a
Distant	n/a
Unstaged	n/a
Overall	67%

n/a - not available

Figures include only invasive cancers


¹ 5-Year Relative Survival Rates for cases diagnosed 2000-2004 based on follow-up of patients through 2005. Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.

Non-Hodgkin Lymphoma

Washington State Residents by Gender

Incidence Rate Trends

Age-adjusted to 2000 US standard population




- For both sexes combined, incidence rates for Non-Hodgkin Lymphoma increased steadily by 1.1% per year during the period from 1992 to 2006.
- For men, incidence rates for Non-Hodgkin Lymphoma increased sharply by 12.4% per year from 1992 to 1994; rates remained level from 1994 to 1998; since 1998 the rates have increased steadily by 1.6% per year.
- For women, incidence rates for Non-Hodgkin Lymphoma increased steadily by 1.3% per year during the period from 1992 to 2006.

Washington State Residents by Gender

Death Rate Trends

Age-adjusted to 2000 US standard population



- For both sexes combined, mortality rates for Non-Hodgkin Lymphoma remained level from 1992 to 2001; since then, however the rates have decreased sharply by 5.1% per year.
- For men, mortality rates for Non-Hodgkin Lymphoma increased steadily by 1.1% per year from 1992 to 2002; since then, however the rates have decreased sharply by 9.0% per year.
- For women, mortality rates for Non-Hodgkin Lymphoma decreased steadily by 2.0% per year during the period from 1992 to 2006.

The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.


Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.

Non-Hodgkin Lymphoma

2004-2006 Washington State Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population




NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.
Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2006 released 1/2009.

Non-Hodgkin Lymphoma

2004-2006 Washington State Residents

Average Annual Death Rates by County with rates in [] and 95% CI's |—|

Age-adjusted to 2000 US standard population



NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.

Oral Cavity and Pharynx

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using primary site ICD-03 codes C00.0-C14.8, excluding histology codes 9140, 9590-9989. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 codes 140.0-149.9 (1992-1998), ICD-10 codes C00-C14 (1999-2006).

2006 Washington State Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


Incidence			
	Male	Female	Total
All Reportable Cancers	493	233	726
2006 Incidence Rates	15.7	6.8	10.9
2005 National Rates	15.3	6.2	10.4

Death			
	Male	Female	Total
Deaths	131	64	195
2006 Death Rates	4.4	1.8	3.0
2005 National Rates	3.8	1.4	2.5

Stage at Diagnosis

2006 Washington State Residents

Total Incidence and Percents



Relative Survival Rates

2005 National 5-year rates¹


In situ	89%
Local	83%
Regional	53%
Distant	30%
Unstaged	49%
Overall	61%

Figures include in situ and invasive cancers

¹ 5-Year Relative Survival Rates for cases diagnosed 2000-2004 based on follow-up of patients through 2005. Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.


Oral Cavity and Pharynx

Washington State Residents by Gender
Incidence Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined, oral cavity and pharyngeal cancer incidence rates decreased steadily by 1% per year during the period from 1992 to 2006.
- For men, oral cavity and pharyngeal cancer incidence rates decreased slightly by 0.8% per year during the period from 1992 to 2006.
- For women, oral cavity and pharyngeal cancer incidence rates decreased steadily by 1.7% per year during the period from 1992 to 2006.

Washington State Residents by Gender
Death Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined and men and women separately, oral cavity and pharynx cancer mortality rates remained level during the period from 1992 to 2006.


The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.
 Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.

Oral Cavity and Pharynx

2004-2006 Washington State Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population




NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.
Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2006 released 1/2009.

Oral Cavity and Pharynx

2004-2006 Washington State Residents

Average Annual Death Rates by County with rates in [] and 95% CI's |—|

Age-adjusted to 2000 US standard population



NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.

Other Endocrine (including thymus)

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using primary site ICD-03 codes C37.9, C74.0-C74.9, C75.0-C75.9, excluding histology codes 9140, 9590-9989. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 codes 164.0, 194 (1992-1998), ICD-10 codes C37, C74-C75 (1999-2006).

2006 Washington State Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


Incidence			
	Male	Female	Total
All Reportable Cancers	24	18	42
2006 Incidence Rates	0.7	0.5	0.6
2005 National Rates	0.7	0.6	0.7

Death			
	Male	Female	Total
Deaths	11	11	22
2006 Death Rates	0.3	0.3	0.3
2005 National Rates	0.3	0.3	0.3

Stage at Diagnosis

2006 Washington State Residents

Total Incidence and Percents



Relative Survival Rates

2005 National 5-year rates¹

In situ	n/a
Local	82%
Regional	70%
Distant	39%
Unstaged	49%
Overall	61%

n/a - not available


n/c - not calculated because the number of cases is less than 5

Figures include in situ and invasive cancers

¹ 5-Year Relative Survival Rates for cases diagnosed 2000-2004 based on follow-up of patients through 2005. Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.


Other Endocrine (including thymus)

Washington State Residents by Gender
Incidence Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined and men and women separately, other endocrine cancer incidence rates remained level during the period from 1992 to 2006.

Washington State Residents by Gender
Death Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined and men and women separately, other endocrine cancer mortality rates remained level during the period from 1992 to 2006.


The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.
 Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.

Other Endocrine (including thymus)

2004-2006 Washington State Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population




NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.
Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2006 released 1/2009.

Other Endocrine (including thymus)

2004-2006 Washington State Residents

Average Annual Death Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population



NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.

Ovary

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using primary site ICD-03 code C56.9, excluding histology codes 9140, 9590-9989. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 code 183.0 (1992-1998), ICD-10 code C56 (1999-2006).

2006 Washington State Female Residents Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


Incidence			
	Male	Female	Total
All Reportable Cancers	n/a	479	n/a
2006 Incidence Rates	n/a	13.7	n/a
2005 National Rates	n/a	12.7	n/a

Death			
	Male	Female	Total
Deaths	n/a	338	n/a
2006 Death Rates	n/a	9.8	n/a
2005 National Rates	n/a	8.6	n/a

Stage at Diagnosis

2006 Washington State Female Residents

Total Incidence and Percents



Relative Survival Rates

2005 National 5-year rates¹

In situ	95%
Local	94%
Regional	74%
Distant	32%
Unstaged	30%
Overall	49%

n/c - not calculated because the number of cases is less than 5

Figures include in situ and invasive cancers


¹ 5-Year Relative Survival Rates for cases diagnosed 2000-2004 based on follow-up of patients through 2005.

n/a Not available for sex-specific sites.

Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.


Ovary

Washington State Female Residents
Incidence Rate Trends
 Age-adjusted to 2000 US standard population



- For women, ovarian cancer incidence rates remained level from 1992 to 2002; since then, however the rates have decreased sharply by 4.2% per year.

Washington State Female Residents
Death Rate Trends
 Age-adjusted to 2000 US standard population



- For women, ovarian cancer mortality rates remained level during the period from 1992 to 2006.

The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.


Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.

Ovary

2004-2006 Washington State Female Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population




NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.
Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2006 released 1/2009.

Ovary

2004-2006 Washington State Female Residents

Average Annual Death Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population



NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.

Pancreas

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using primary site ICD-0 codes C25.0-C25.9, excluding histology codes 9140, 9590-9989. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 codes 157.0-157.9 (1992-1998), ICD-10 code C25 (1999-2006).

2006 Washington State Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


Incidence			
	Male	Female	Total
All Reportable Cancers	380	359	739
2006 Incidence Rates	13.9	10.3	11.9
2005 National Rates	12.9	10.4	11.5

Death			
	Male	Female	Total
Deaths	326	324	650
2006 Death Rates	11.9	9.1	10.3
2005 National Rates	12.4	9.4	10.8

Stage at Diagnosis

2006 Washington State Residents

Total Incidence and Percents



Relative Survival Rates

2005 National 5-year rates¹

In situ	90%
Local	21%
Regional	8%
Distant	2%
Unstaged	4%
Overall	5%


n/c - not calculated because the number of cases is less than 5

Figures include in situ and invasive cancers

¹ 5-Year Relative Survival Rates for cases diagnosed 2000-2004 based on follow-up of patients through 2005. Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.


Pancreas

Washington State Residents by Gender
Incidence Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined, pancreatic cancer incidence rates increased slightly by 0.7% per year during the period from 1992 to 2006.
- For men, pancreatic cancer incidence rates increased steadily by 1% per year during the period from 1992 to 2006.
- For women, pancreatic cancer incidence rates remained level during the period from 1992 to 2006.

Washington State Residents by Gender
Death Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined and men and women separately, pancreatic cancer mortality rates remained level during the period from 1992 to 2006.


The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.
 Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.

Pancreas

2004-2006 Washington State Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population




NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.
Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2006 released 1/2009.

Pancreas

2004-2006 Washington State Residents

Average Annual Death Rates by County with rates in [] and 95% CI's |—|

Age-adjusted to 2000 US standard population



NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.

Prostate

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using primary site ICD-03 code C61.9, excluding histology codes 9140, 9590-9989. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 code 185 (1992-1998), ICD-10 code C61 (1999-2006).

2006 Washington State Male Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


Incidence			
	Male	Female	Total
All Reportable Cancers	4,845	n/a	n/a
2006 Incidence Rates	166.9	n/a	n/a
2005 National Rates	144.2	n/a	n/a

Death			
	Male	Female	Total
Deaths	610	n/a	n/a
2006 Death Rates	25.1	n/a	n/a
2005 National Rates	24.6	n/a	n/a

Stage at Diagnosis

2006 Washington State Male Residents

Total Incidence and Percents



Relative Survival Rates

2005 National 5-year rates¹

In situ	90%
Local	100%
Regional	100%
Distant	29%
Unstaged	77%
Overall	100%

n/c - not calculated because the number of cases is less than 5

Figures include in situ and invasive cancers

¹ 5-Year Relative Survival Rates for cases diagnosed 2000-2004 based on follow-up of patients through 2005.

n/a Not available for sex-specific sites.

Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.

Prostate

Washington State Male Residents

Incidence Rate Trends

Age-adjusted to 2000 US standard population




- For men, prostate cancer incidence rates decreased sharply by 12.7% per year from 1992 to 1995; rates remained level from 1995 to 1999; since 1999 the rates have decreased steadily by 2.0% per year.

Washington State Male Residents

Death Rate Trends

Age-adjusted to 2000 US standard population



- For men, prostate cancer mortality rates decreased sharply by 4.9% per year from 1992 to 1998; since 1998 the rates have decreased steadily by 2.1% per year.

The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.


Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.

Prostate

2004-2006 Washington State Male Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population




NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.
Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2006 released 1/2009.

Prostate

2004-2006 Washington State Male Residents

Average Annual Death Rates by County with rates in [] and 95% CI 's |

Age-adjusted to 2000 US standard population



NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.

Stomach

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using primary site ICD-03 codes C16.0-C16.9, excluding histology codes 9140, 9590-9989. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 codes 151.0-151.9 (1992-1998), ICD-10 code C16 (1999-2006).

2006 Washington State Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


Incidence			
	Male	Female	Total
All Reportable Cancers	258	142	400
2006 Incidence Rates	9.3	4.2	6.4
2005 National Rates	10.9	5.3	7.7

Death			
	Male	Female	Total
Deaths	129	86	215
2006 Death Rates	4.8	2.4	3.4
2005 National Rates	5.2	2.7	3.8

Stage at Diagnosis

2006 Washington State Residents

Total Incidence and Percents



Relative Survival Rates

2005 National 5-year rates¹

In situ	74%
Local	60%
Regional	26%
Distant	4%
Unstaged	16%
Overall	26%


n/c - not calculated because the number of cases is less than 5

Figures include in situ and invasive cancers

¹ 5-Year Relative Survival Rates for cases diagnosed 2000-2004 based on follow-up of patients through 2005. Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.


Stomach

Washington State Residents by Gender
Incidence Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined, stomach cancer incidence rates decreased steadily by 1.0% per year during the period from 1992 to 2006.
- For men, stomach cancer incidence rates decreased steadily by 1.5% per year during the period from 1992 to 2006.
- For women, stomach cancer incidence rates remained level during the period from 1992 to 2006.

Washington State Residents by Gender
Death Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined, stomach cancer mortality rates decreased steadily by 3.1% per year during the period from 1992 to 2006.
- For men, stomach cancer mortality rates decreased steadily by 3.9% per year during the period from 1992 to 2006.
- For women, stomach cancer mortality rates decreased steadily by 2.0% per year during the period from 1992 to 2006.

The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.


Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.

Stomach

2004-2006 Washington State Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population




NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.
Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2006 released 1/2009.

Stomach

2004-2006 Washington State Residents

Average Annual Death Rates by County with rates in [] and 95% CI's |—|

Age-adjusted to 2000 US standard population



NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.

Testis

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using primary site ICD-03 codes C62.0-C62.9, excluding histology codes 9140, 9590-9989. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 codes 186.0-186.9 (1992-1998), ICD-10 code C62 (1999-2006).

2006 Washington State Male Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


Incidence			
	Male	Female	Total
All Reportable Cancers	233	n/a	n/a
2006 Incidence Rates	7.1	n/a	n/a
2005 National Rates	5.4	n/a	n/a

Death			
	Male	Female	Total
Deaths	6	n/a	n/a
2006 Death Rates	0.2	n/a	n/a
2005 National Rates	0.2	n/a	n/a

Stage at Diagnosis

2006 Washington State Male Residents

Total Incidence and Percents



Relative Survival Rates	
2005 National 5-year rates ¹	
In situ	100%
Local	99%
Regional	96%
Distant	72%
Unstaged	85%
Overall	96%

n/c - not calculated because the number of cases is less than 5

Figures include in situ and invasive cancers

¹ 5-Year Relative Survival Rates for cases diagnosed 2000-2004 based on follow-up of patients through 2005.

n/a Not available for sex-specific sites.

Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.


Testis

Washington State Male Residents
Incidence Rate Trends
 Age-adjusted to 2000 US standard population



- For men, testicular cancer incidence rates increased slightly by 0.7% per year during the period from 1992 to 2006.

Washington State Male Residents
Death Rate Trends
 Age-adjusted to 2000 US standard population



- For men, testicular cancer mortality rates remained level during the period from 1992 to 2006.

The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.


Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.

Testis

2004-2006 Washington State Male Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population




NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.
Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2006 released 1/2009.

Testis

2004-2006 Washington State Male Residents

Average Annual Death Rates by County with rates in [] and 95% CI 's |

Age-adjusted to 2000 US standard population



NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.

Thyroid

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using primary site ICD-03 code C73.9, excluding histology codes 9140, 9590-9989. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 code 193 (1992-1998), ICD-10 code C73 (1999-2006).

2006 Washington State Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


Incidence			
	Male	Female	Total
All Reportable Cancers	206	548	754
2006 Incidence Rates	6.5	16.5	11.5
2005 National Rates	5.3	15.2	10.3

Death			
	Male	Female	Total
Deaths	12	13	25
2006 Death Rates	0.4	0.4	0.4
2005 National Rates	0.5	0.5	0.5

Stage at Diagnosis

2006 Washington State Residents

Total Incidence and Percents



Relative Survival Rates

2005 National 5-year rates¹

In situ	n/a
Local	100%
Regional	97%
Distant	58%
Unstaged	90%
Overall	97%


n/a - not available

Figures include only invasive cancers

¹ 5-Year Relative Survival Rates for cases diagnosed 2000-2004 based on follow-up of patients through 2005. Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.


Thyroid

Washington State Residents by Gender
Incidence Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined, thyroid cancer incidence rates increased sharply by 5.4% per year during the period from 1992 to 2006.
- For men, thyroid cancer incidence rates increased steadily by 3.7% per year from 1992 to 2004; since then, however the rates have remained level.
- For women, thyroid cancer incidence rates increased sharply by 5.5% per year during the period from 1992 to 2006.

Washington State Residents by Gender
Death Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined, men and women separately, thyroid cancer mortality rates remained level during the period from 1992 to 2006.


The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.
 Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.

Thyroid

2004-2006 Washington State Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population




NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.
Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2006 released 1/2009.

Thyroid

2004-2006 Washington State Residents

Average Annual Death Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population



NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.

Vulva (Female)

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using primary site ICD-03 code C51.0-C51.9, excluding histology codes 9140, 9590-9989. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 code 184.1-184.4 (1992-1998), ICD-10 code C51 (1999-2006).

2006 Washington State Female Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


Incidence			
	Male	Female	Total
All Reportable Cancers	n/a	264	n/a
2006 Incidence Rates	n/a	7.9	n/a
2005 National Rates	n/a	5.8	n/a

Death			
	Male	Female	Total
Deaths	n/a	20	n/a
2006 Death Rates	n/a	0.6	n/a
2005 National Rates	n/a	0.4	n/a

Stage at Diagnosis

2006 Washington State Female Residents

Total Incidence and Percents



Relative Survival Rates

2005 National 5-year rates¹

In situ	98%
Local	91%
Regional	56%
Distant	14%
Unstaged	54%
Overall	90%

n/c - not calculated because the number of cases is less than 5

Figures include in situ and invasive cancers


¹ 5-Year Relative Survival Rates for cases diagnosed 2000-2004 based on follow-up of patients through 2005.

n/a Not available for sex-specific sites.

Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.


Vulva (Female)

Washington State Female Residents
Incidence Rate Trends
 Age-adjusted to 2000 US standard population



- For women, vulva cancer incidence rates remained level during the period from 1992 to 2006.

Washington State Female Residents
Death Rate Trends
 Age-adjusted to 2000 US standard population



- For women, vulva cancer mortality rates remained level during the period from 1992 to 2006.


The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.
 Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2006 released in January 2009.

Vulva (Female)

2004-2006 Washington State Female Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population




NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.
Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2006 released 1/2009.

Vulva (Female)

2004-2006 Washington State Female Residents

Average Annual Death Rates by County with rates in [] and 95% CI's |—|

Age-adjusted to 2000 US standard population



NOTE: Rates and confidence intervals are not computed when the 2004-2006 total number of new cases is less than 5.