



Cancer By Site

Percent Distribution of Cancer

2007 Total Cancer Incidence of Washington State Residents




2007 Total Cancer Mortality of Washington State Residents




Percent Distribution of Cancer

2007 Male Cancer Incidence of Washington State Residents




2007 Male Cancer Mortality of Washington State Residents




Percent Distribution of Cancer

2007 Female Cancer Incidence of Washington State Residents



2007 Female Cancer Mortality of Washington State Residents



All Sites Combined

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using ICD-03 codes C00.0 - C80.9, all histology codes. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 codes 140-208, 238.6 (1992-1998), ICD-10 codes C00-C97 (1999-2007).

2007 Washington State Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


| Incidence | | | |
|------------------------|--------|--------|--------|
| | Male | Female | Total |
| All Reportable Cancers | 16,572 | 16,501 | 33,077 |
| 2007 Incidence Rates | 560.2 | 470.8 | 505.5 |
| 2006 National Rates | 554.0 | 455.3 | 493.6 |

| Death | | | |
|---------------------|-------|--------|--------|
| | Male | Female | Total |
| Deaths | 5,920 | 5,605 | 11,525 |
| 2007 Death Rates | 213.4 | 156.1 | 179.5 |
| 2006 National Rates | 221.1 | 153.7 | 181.1 |

Stage at Diagnosis

2007 Washington State Residents

Total Incidence and Percents



Relative Survival Rates

2006 National 5-year rates¹

| | |
|----------|-----|
| In situ | n/a |
| Local | n/a |
| Regional | n/a |
| Distant | n/a |
| Unstaged | n/a |
| Overall | 69% |


n/a = not available

Figures include in situ and invasive cancers

¹ 5-Year Relative Survival Rates for cases diagnosed 2001-2005 based on follow-up of patients through 2006. Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.


All Sites Combined

**Washington State Residents by Gender
Incidence Rate Trends**
Age-adjusted to 2000 US standard population



- For both sexes combined, cancer incidence rates for all cancer sites combined increased slightly by 0.8% per year from 1992 to 2001; since then, however the rates have decreased steadily by 1.2 % per year.
- For men, cancer incidence rates for all cancer sites combined remained level from 1992 to 1999; since then, however the rates have decreased steadily by 1.0% per year.
- For women, cancer incidence rates for all cancer sites combined increased steadily by 2.0% per year from 1992 to 1998; since then, however the rates have remained level.

**Washington State Residents by Gender
Death Rate Trends**
Age-adjusted to 2000 US standard population



- For both sexes combined, cancer mortality rates for all cancer sites combined decreased steadily by 1.1% per year during the period from 1992 to 2007.
- For men, cancer mortality rates for all cancer sites combined decreased steadily by 1.5% per year during the period from 1992 to 2007.
- For women, cancer mortality rates for all cancer sites combined decreased slightly by 0.9% per year during the period from 1992 to 2007.

The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.


Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.

All Sites Combined

2005-2007 Washington State Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population




Age-adjusted rate per 100,000
 NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.
 Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2007 released 1/2010,

All Sites Combined

2005-2007 Washington State Residents

Average Annual Death Rates by County with rates in [] and 95% CI's |—|

Age-adjusted to 2000 US standard population



Age-adjusted rate per 100,000

NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.

Bladder

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using primary site ICD-03 codes C67.0-C67.9, excluding histology codes 9140, 9590-9989. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 codes 188.0-188.9 (1992-1998), ICD-10 code C67 (1999-2007).

2007 Washington State Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


| Incidence | | | |
|------------------------|-------|--------|-------|
| | Male | Female | Total |
| All Reportable Cancers | 1,009 | 322 | 1,331 |
| 2007 Incidence Rates | 36.1 | 9.1 | 20.8 |
| 2006 National Rates | 36.0 | 8.7 | 20.3 |

| Death | | | |
|---------------------|------|--------|-------|
| | Male | Female | Total |
| Deaths | 232 | 84 | 316 |
| 2007 Death Rates | 8.7 | 2.2 | 4.9 |
| 2006 National Rates | 7.6 | 2.2 | 4.3 |

Stage at Diagnosis

2007 Washington State Residents

Total Incidence and Percents



Relative Survival Rates

2006 National 5-year rates¹

| | |
|----------|-----|
| In situ | n/a |
| Local | 93% |
| Regional | 44% |
| Distant | 6% |
| Unstaged | 54% |
| Overall | 80% |


n/a = not available

Figures include in situ and invasive cancers

¹ 5-Year Relative Survival Rates for cases diagnosed 2001-2005 based on follow-up of patients through 2006. Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.


Bladder

**Washington State Residents by Gender
Incidence Rate Trends**
Age-adjusted to 2000 US standard population



- For both sexes combined, bladder cancer incidence rates increased slightly by 0.6% per year from 1992 to 2005; since then, however the rates have decreased sharply by 8.0% per year.
- For men, bladder cancer incidence rates increased steadily by 2.4% per year from 1992 to 1998; rates remained level from 1998 to 2005; since 2005 the rates have decreased sharply by 8.5% per year.
- For women, bladder cancer incidence rates remained level from 1992 to 2002; since then, however the rates have decreased sharply by 3.5% per year.

**Washington State Residents by Gender
Death Rate Trends**
Age-adjusted to 2000 US standard population



- For both sexes combined and men and women separately, bladder cancer mortality rates remained level during the period from 1992 to 2007.

The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.


Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.

Bladder

2005-2007 Washington State Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population




Age-adjusted rate per 100,000
 NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.
 Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2007 released 1/2010.

Bladder

2005-2007 Washington State Residents

Average Annual Death Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population



Age-adjusted rate per 100,000
 NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.

Brain

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using primary site ICD-03 codes C71.0-C71.9, excluding histology codes 9530-9539 and 9590-9989. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 codes 191.0-191.9 (1992-1998), ICD-10 code C71 (1999-2007).

2007 Washington State Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


| Incidence | | | |
|------------------------|------|--------|-------|
| | Male | Female | Total |
| All Reportable Cancers | 231 | 190 | 421 |
| 2007 Incidence Rates | 7.5 | 5.5 | 6.4 |
| 2006 National Rates | 7.0 | 4.8 | 5.8 |

| Death | | | |
|---------------------|------|--------|-------|
| | Male | Female | Total |
| Deaths | 172 | 137 | 309 |
| 2007 Death Rates | 5.6 | 3.9 | 4.7 |
| 2006 National Rates | 5.1 | 3.4 | 4.2 |

Stage at Diagnosis

2007 Washington State Residents

Total Incidence and Percents



Relative Survival Rates

2006 National 5-year rates¹

| | |
|----------|-----|
| In situ | n/a |
| Local | n/a |
| Regional | n/a |
| Distant | n/a |
| Unstaged | n/a |
| Overall | 31% |

n/a = not available


n/c Number of cases <= 5

Figures include only invasive cancers

¹ 5-Year Relative Survival Rates for cases diagnosed 2001-2005 based on follow-up of patients through 2006. Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.


Brain

**Washington State Residents by Gender
Incidence Rate Trends**
Age-adjusted to 2000 US standard population



- For both sexes combined, brain cancer incidence rates increased steadily by 2.7% per year from 1992 to 1999; since then, however the rates have decreased steadily by 1.7% per year.
- For men and women separately, brain cancer incidence rates remained level during the period from 1992 to 2007.

**Washington State Residents by Gender
Death Rate Trends**
Age-adjusted to 2000 US standard population



- For both sexes combined, brain cancer mortality rates decreased slightly by 0.8% per year during the period from 1992 to 2007.
- For men, brain cancer mortality rates decreased steadily by 1.0% per year during the period from 1992 to 2007.
- For women, brain cancer mortality rates remained level during the period from 1992 to 2007.


The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.

Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.

Brain

2005-2007 Washington State Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |
 Age-adjusted to 2000 US standard population




Age-adjusted rate per 100,000
 NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.
 Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2007 released 1/2010.

Brain

2005-2007 Washington State Residents

Average Annual Death Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population



Age-adjusted rate per 100,000
 NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.

Breast (Female)

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using primary site ICD-03 codes C50.0-C50.9, excluding histology codes 9140, 9590-9989. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 codes 174.0-174.9 (1992-1998), ICD-10 codes C50 (1999-2007).

2007 Washington State Female Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


| Incidence | | | |
|------------------------|------|--------|-------|
| | Male | Female | Total |
| All Reportable Cancers | n/a | 5,563 | n/a |
| 2007 Incidence Rates | n/a | 158.4 | n/a |
| 2006 National Rates | n/a | 150.9 | n/a |

| Death | | | |
|---------------------|------|--------|-------|
| | Male | Female | Total |
| Deaths | n/a | 795 | n/a |
| 2007 Death Rates | n/a | 21.9 | n/a |
| 2006 National Rates | n/a | 23.4 | n/a |

Stage at Diagnosis

2007 Washington State Female Residents

Total Incidence and Percents



Relative Survival Rates

2006 National 5-year rates¹

| | |
|----------|------|
| In situ | 100% |
| Local | 98% |
| Regional | 85% |
| Distant | 29% |
| Unstaged | 53% |
| Overall | 92% |

Figures include in situ and invasive cancers


¹ 5-Year Relative Survival Rates for cases diagnosed 2001-2005 based on follow-up of patients through 2006.

n/a Not available for sex-specific sites.

Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.


Breast (Female)

Washington State Female Residents
Incidence Rate Trends
 Age-adjusted to 2000 US standard population



- For women, breast cancer incidence rates increased steadily by 2.9% per year from 1992 to 1999; since then, however the rates have decreased steadily by 2.1% per year.

Washington State Female Residents
Death Rate Trends
 Age-adjusted to 2000 US standard population



- For women, breast cancer mortality rates decreased steadily by 2.3% per year during the period from 1992 to 2007.

The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.


Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry- Incidence data for diagnosis years 1992 to 2007 released in January 2010.

Breast (Female)

2005-2007 Washington State Female Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population




Age-adjusted rate per 100,000
 NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.
 Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2007 released 1/2010.

Breast (Female)

2005-2007 Washington State Female Residents

Average Annual Death Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population



Age-adjusted rate per 100,000

NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.

Cervix (Uterine)

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using primary site ICD-03 codes C53.0-C53.9, excluding histology codes 9140, 9590-9989 and excluding in situ. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 codes 180.0-180.9 (1992-1998), ICD-10 code C53 (1999-2007).

2007 Washington State Female Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


| Incidence | | | |
|------------------------|------|--------|-------|
| | Male | Female | Total |
| All Reportable Cancers | n/a | 196 | n/a |
| 2007 Incidence Rates | n/a | 6.0 | n/a |
| 2006 National Rates | n/a | 8.0 | n/a |

| Death | | | |
|---------------------|------|--------|-------|
| | Male | Female | Total |
| Deaths | n/a | 79 | n/a |
| 2007 Death Rates | n/a | 2.2 | n/a |
| 2006 National Rates | n/a | 2.4 | n/a |

Stage at Diagnosis

2007 Washington State Female Residents

Total Incidence and Percents



Relative Survival Rates

2006 National 5-year rates¹

| | |
|----------|-----|
| In situ | n/a |
| Local | 91% |
| Regional | 56% |
| Distant | 16% |
| Unstaged | 55% |
| Overall | 69% |

n/a = not available

Figures include only invasive cancers


¹ 5-Year Relative Survival Rates for cases diagnosed 2001-2005 based on follow-up of patients through 2006.

n/a Not available for sex-specific sites.

Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.


Cervix (Uterine)

**Washington State Female Residents
Incidence Rate Trends**
Age-adjusted to 2000 US standard population



- For women, cervical cancer incidence rates decreased steadily by 2.4% per year during the period from 1992 to 2007.

**Washington State Female Residents
Death Rate Trends**
Age-adjusted to 2000 US standard population



- For women, cervical cancer mortality rates decreased steadily by 2.0% per year during the period from 1992 to 2007.

The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.


Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry- Incidence data for diagnosis years 1992 to 2007 released in January 2010.

Cervix (Uterine)

2005-2007 Washington State Female Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population




Age-adjusted rate per 100,000
 NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.
 Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2007 released 1/2010.

Cervix (Uterine)

2005-2007 Washington State Female Residents

Average Annual Death Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population



Age-adjusted rate per 100,000
 NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.

Colorectal

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using primary site ICD-03 codes C18.0-C18.9, C19.9, C20.9, C26.0, excluding histology codes 9140, 9590-9989. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 codes 153.0-154.1, 159.0 (1992-1998), ICD-10 codes C18-C20, C26.0 (1999-2007).

2007 Washington State Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


| Incidence | | | |
|------------------------|-------|--------|-------|
| | Male | Female | Total |
| All Reportable Cancers | 1,392 | 1,244 | 2,636 |
| 2007 Incidence Rates | 47.4 | 34.7 | 40.6 |
| 2006 National Rates | 55.8 | 42.6 | 48.5 |

| Death | | | |
|---------------------|------|--------|-------|
| | Male | Female | Total |
| Deaths | 498 | 474 | 972 |
| 2007 Death Rates | 17.6 | 12.5 | 14.9 |
| 2006 National Rates | 20.5 | 14.5 | 17.1 |

Stage at Diagnosis

2007 Washington State Residents

Total Incidence and Percents



Relative Survival Rates

2006 National 5-year rates¹


| | |
|----------|-----|
| In situ | 96% |
| Local | 90% |
| Regional | 70% |
| Distant | 12% |
| Unstaged | 38% |
| Overall | 67% |

Figures include in situ and invasive cancers

¹ 5-Year Relative Survival Rates for cases diagnosed 2001-2005 based on follow-up of patients through 2006. Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.


Colorectal

**Washington State Residents by Gender
Incidence Rate Trends**
Age-adjusted to 2000 US standard population



- For both sexes combined, colorectal cancer incidence rates remained level from 1992 to 1999; since then, however the rates have decreased steadily by 3.8% per year.
- For men, colorectal cancer incidence rates remained level from 1992 to 1998; since then, however the rates have decreased steadily by 3.8% per year.
- For women, colorectal cancer incidence rates remained level from 1992 to 1998; since then, however the rates have decreased steadily by 3.3% per year.

**Washington State Residents by Gender
Death Rate Trends**
Age-adjusted to 2000 US standard population



- For both sexes combined, colorectal cancer mortality rates decreased steadily by 2.6% per year during the period from 1992 to 2007.
- For men, colorectal cancer mortality rates decreased steadily by 2.8% per year during the period from 1992 to 2007.
- For women, colorectal cancer mortality rates decreased steadily by 2.7% per year during the period from 1992 to 2007.

The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.


Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.

Colorectal

2005-2007 Washington State Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's

Age-adjusted to 2000 US standard population




Age-adjusted rate per 100,000
 NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.
 Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2007 released

Colorectal

2005-2007 Washington State Residents

Average Annual Death Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population



Age-adjusted rate per 100,000
 NOTE: Rates and confidence intervals are not computed when the
 2005-2007 total number of new cases <= 5

Endometrium

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using primary site ICD-03 codes C54.0-C54.9, excluding histology codes 9140, 9590-9989. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 codes 182.0-182.1, 182.8 (1992-1998), ICD-10 codes C54 (1999-2007).

2007 Washington State Female Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


| Incidence | | | |
|------------------------|------|--------|-------|
| | Male | Female | Total |
| All Reportable Cancers | n/a | 790 | n/a |
| 2007 Incidence Rates | n/a | 22.0 | n/a |
| 2006 National Rates | n/a | 23.2 | n/a |

| Death | | | |
|---------------------|------|--------|-------|
| | Male | Female | Total |
| Deaths | n/a | 70 | n/a |
| 2007 Death Rates | n/a | 1.9 | n/a |
| 2006 National Rates | n/a | 2.0 | n/a |

Stage at Diagnosis

2007 Washington State Female Residents

Total Incidence and Percents



Relative Survival Rates

2006 National 5-year rates¹

| | |
|----------|-----|
| In situ | 99% |
| Local | 96% |
| Regional | 69% |
| Distant | 25% |
| Unstaged | 58% |
| Overall | 84% |

n/c Number of cases <= 5

Figures include in situ and invasive cancers


¹ 5-Year Relative Survival Rates for cases diagnosed 2001-2005 based on follow-up of patients through 2006.

n/a Not available for sex-specific sites.

Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.


Endometrium

Washington State Female Residents
Incidence Rate Trends
 Age-adjusted to 2000 US standard population



- For women, endometrial cancer incidence rates decreased steadily by 1.1% per year during the period from 1992 to 2007.

Washington State Female Residents
Death Rate Trends
 Age-adjusted to 2000 US standard population



- For women, endometrial cancer mortality rates remained level during the period from 1992 to 2007.

The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.


Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry- Incidence data for diagnosis years 1992 to 2007 released in January 2010.

Endometrium

2005-2007 Washington State Female Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population



Age-adjusted rate per 100,000

NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.


Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2007 released 1/2010.

Endometrium

2005-2007 Washington State Female Residents

Average Annual Death Rates by County with rates in [] and 95% CI's

Age-adjusted to 2000 US standard population



Age-adjusted rate per 100,000
 NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.

Esophagus

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using primary site ICD-03 codes C15.0-C15.9, excluding histology codes 9140, 9590-9989. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 codes 150.0-150.9 (1992-1998), ICD-10 code C15 (1999-2007).

2007 Washington State Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


| Incidence | | | |
|------------------------|------|--------|-------|
| | Male | Female | Total |
| All Reportable Cancers | 250 | 67 | 317 |
| 2007 Incidence Rates | 8.6 | 1.9 | 4.9 |
| 2006 National Rates | 7.6 | 2.0 | 4.5 |

| Death | | | |
|---------------------|------|--------|-------|
| | Male | Female | Total |
| Deaths | 243 | 77 | 320 |
| 2007 Death Rates | 8.5 | 2.2 | 5.0 |
| 2006 National Rates | 7.9 | 1.7 | 4.4 |

Stage at Diagnosis

2007 Washington State Residents

Total Incidence and Percents



Relative Survival Rates

2006 National 5-year rates¹

| | |
|----------|-----|
| In situ | 74% |
| Local | 38% |
| Regional | 18% |
| Distant | 3% |
| Unstaged | 12% |
| Overall | 18% |


n/c Number of cases <= 5

Figures include in situ and invasive cancers

¹ 5-Year Relative Survival Rates for cases diagnosed 2001-2005 based on follow-up of patients through 2006. Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.


Esophagus

**Washington State Residents by Gender
Incidence Rate Trends**
Age-adjusted to 2000 US standard population



- For both sexes combined, esophageal cancer incidence rates increased steadily by 3.0% per year from 1992 to 2000; since then, however the rates have decreased steadily by 3.1% per year.
- For men, esophageal cancer incidence rates increased sharply by 3.9% per year from 1992 to 1999; since then, however the rates have decreased steadily by 2.9% per year.
- For women, esophageal cancer incidence rates remained level during the period from 1992 to 2007.

**Washington State Residents by Gender
Death Rate Trends**
Age-adjusted to 2000 US standard population



- For both sexes combined, esophageal cancer mortality rates increased slightly by 0.8% per year during the period from 1992 to 2007.
- For men, esophageal cancer mortality rates increased slightly by 0.9% per year during the period from 1992 to 2007.
- For women, esophageal cancer mortality rates remained level during the period from 1992 to 2007.

The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.


Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.

Esophagus

2005-2007 Washington State Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population




Age-adjusted rate per 100,000
 NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.
 Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2007 released 1/2010.

Esophagus

2005-2007 Washington State Residents

Average Annual Death Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population



Age-adjusted rate per 100,000
 NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.

Hodgkin Lymphoma

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using all primary sites with histology codes 9650-9667. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 codes 201.0-201.9 (1992-1998), ICD-10 code C81 (1999-2007).

2007 Washington State Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


| Incidence | | | |
|------------------------|------|--------|-------|
| | Male | Female | Total |
| All Reportable Cancers | 101 | 84 | 185 |
| 2007 Incidence Rates | 3.1 | 2.6 | 2.8 |
| 2006 National Rates | 3.2 | 2.4 | 2.8 |

| Death | | | |
|---------------------|------|--------|-------|
| | Male | Female | Total |
| Deaths | 15 | 12 | 27 |
| 2007 Death Rates | 0.5 | 0.3 | 0.4 |
| 2006 National Rates | 0.6 | 0.3 | 0.4 |

Stage at Diagnosis

2007 Washington State Residents

Total Incidence and Percents



Relative Survival Rates

2006 National 5-year rates¹

| | |
|----------|-----|
| In situ | n/a |
| Local | n/a |
| Regional | n/a |
| Distant | n/a |
| Unstaged | n/a |
| Overall | 85% |


n/a = not available

Figures include only invasive cancers

¹ 5-Year Relative Survival Rates for cases diagnosed 2001-2005 based on follow-up of patients through 2006. Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.


Hodgkin Lymphoma

**Washington State Residents by Gender
Incidence Rate Trends**
Age-adjusted to 2000 US standard population



- For both sexes combined and men and women separately, incidence rates for Hodgkin Lymphoma remained level during the period from 1992 to 2007.

**Washington State Residents by Gender
Death Rate Trends**
Age-adjusted to 2000 US standard population



- For both sexes combined, mortality rates for Hodgkin Lymphoma decreased steadily by 2.4% per year during the period from 1992 to 2007.
- For men and women separately, mortality rates for Hodgkin Lymphoma remained level during the period from 1992 to 2007.

The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.


Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.

Hodgkin Lymphoma

2005-2007 Washington State Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population




Age-adjusted rate per 100,000
 NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.
 Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2007 released 1/2010.

Hodgkin Lymphoma

2005-2007 Washington State Residents

Average Annual Death Rates by County with rates in [] and 95% CI's []

Age-adjusted to 2000 US standard population



Age-adjusted rate per 100,000
 NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.

Kidney and Renal Pelvis

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using primary site ICD-03 codes C64.9, C65.9, excluding histology codes 9140, 9590-9989. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 codes 189.0, 189.1 (1992-1998), ICD-10 codes C64-C65 (1999-2007).

2007 Washington State Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


| Incidence | | | |
|------------------------|------|--------|-------|
| | Male | Female | Total |
| All Reportable Cancers | 651 | 400 | 1,051 |
| 2007 Incidence Rates | 21.4 | 11.3 | 15.9 |
| 2006 National Rates | 19.3 | 10.3 | 14.4 |

| Death | | | |
|---------------------|------|--------|-------|
| | Male | Female | Total |
| Deaths | 167 | 103 | 270 |
| 2007 Death Rates | 5.8 | 2.8 | 4.1 |
| 2006 National Rates | 5.7 | 2.6 | 4.0 |

Stage at Diagnosis

2007 Washington State Residents

Total Incidence and Percents



Relative Survival Rates

2006 National 5-year rates¹


| | |
|----------|-----|
| In situ | 93% |
| Local | 91% |
| Regional | 63% |
| Distant | 11% |
| Unstaged | 38% |
| Overall | 70% |

Figures include in situ and invasive cancers

¹ 5-Year Relative Survival Rates for cases diagnosed 2001-2005 based on follow-up of patients through 2006. Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.


Kidney and Renal Pelvis

Washington State Residents by Gender
Incidence Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined, kidney and renal pelvis cancer incidence rates increased steadily by 2.6% per year during the period from 1992 to 2007.
- For men, kidney and renal pelvis cancer incidence rates increased steadily by 2.4% per year during the period from 1992 to 2007.
- For women, kidney and renal pelvis cancer incidence rates remained level from 1992 to 1994; since then, however the rates have increased steadily by 3.5% per year.

Washington State Residents by Gender
Death Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined and men and women separately, kidney and renal pelvis cancer mortality rates remained level during the period from 1992 to 2007.

The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.


Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.

Kidney and Renal Pelvis

2005-2007 Washington State Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population




Age-adjusted rate per 100,000
 NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.
 Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2007 released 1/2010.

Kidney and Renal Pelvis

2005-2007 Washington State Residents

Average Annual Death Rates by County with rates in [] and 95% CI's |—|

Age-adjusted to 2000 US standard population



NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.

Larynx

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using primary site ICD-03 codes C32.0-C32.9, excluding histology codes 9140, 9590-9989. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 codes 161.0-161.9 (1992-1998), ICD-10 code C32 (1999-2007).

2007 Washington State Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


| Incidence | | | |
|------------------------|------|--------|-------|
| | Male | Female | Total |
| All Reportable Cancers | 161 | 31 | 192 |
| 2007 Incidence Rates | 5.4 | 0.9 | 2.9 |
| 2006 National Rates | 6.5 | 1.3 | 3.6 |

| Death | | | |
|---------------------|------|--------|-------|
| | Male | Female | Total |
| Deaths | 40 | 17 | 57 |
| 2007 Death Rates | 1.4 | 0.5 | 0.9 |
| 2006 National Rates | 2.2 | 0.5 | 1.2 |

Stage at Diagnosis

2007 Washington State Residents

Total Incidence and Percents



Relative Survival Rates

2006 National 5-year rates¹

| | |
|----------|-----|
| In situ | n/a |
| Local | n/a |
| Regional | n/a |
| Distant | n/a |
| Unstaged | n/a |
| Overall | 63% |

n/a = not available


n/c Number of cases ≤ 5

Figures include in situ and invasive cancers

¹ 5-Year Relative Survival Rates for cases diagnosed 2001-2005 based on follow-up of patients through 2006. Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.


Larynx

Washington State Residents by Gender Incidence Rate Trends Age-adjusted to 2000 US standard population



- For both sexes combined, laryngeal cancer incidence rates decreased steadily by 2.9% per year during the period from 1992 to 2007.
- For men, laryngeal cancer incidence rates decreased steadily by 2.9% per year during the period from 1992 to 2007.
- For women, laryngeal cancer incidence rates decreased steadily by 2.6% per year during the period from 1992 to 2007.

Washington State Residents by Gender Death Rate Trends Age-adjusted to 2000 US standard population



- For both sexes combined, laryngeal cancer mortality rates decreased steadily by 1.8% per year during the period from 1992 to 2007.
- For men, laryngeal cancer mortality rates decreased steadily by 2.4% per year during the period from 1992 to 2007.
- For women, laryngeal cancer mortality rates remained level during the period from 1992 to 2007.

The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.


Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.

Larynx

2005-2007 Washington State Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population




Age-adjusted rate per 100,000
 NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.
 Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2007 released 1/2010.

Larynx

2005-2007 Washington State Residents

Average Annual Death Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population



Age-adjusted rate per 100,000
 NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.

Leukemia

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using all sites with histology 9733, 9742, 9800-9820, 9826, 9831-9948, 9963,9964; for sites C420, C421 and C424 only, histology 9823 or 9827. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 codes 204.0-208.9 (1992-1998), ICD-10 codes C90.1, C91-C95 (1999-2007).

2007 Washington State Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


| Incidence | | | |
|------------------------|------|--------|-------|
| | Male | Female | Total |
| All Reportable Cancers | 501 | 341 | 842 |
| 2007 Incidence Rates | 17.4 | 9.8 | 13.1 |
| 2006 National Rates | 14.5 | 9.2 | 11.5 |

| Death | | | |
|---------------------|------|--------|-------|
| | Male | Female | Total |
| Deaths | 288 | 188 | 476 |
| 2007 Death Rates | 10.9 | 5.2 | 7.5 |
| 2006 National Rates | 9.6 | 5.4 | 7.2 |

Stage at Diagnosis

2007 Washington State Residents

Total Incidence and Percents



Relative Survival Rates

2006 National 5-year rates¹

| | |
|----------|-----|
| In situ | n/a |
| Local | n/a |
| Regional | n/a |
| Distant | 54% |
| Unstaged | n/a |
| Overall | 54% |

n/a = not available


n/c Number of cases <= 5

Figures include only invasive cancers

¹ 5-Year Relative Survival Rates for cases diagnosed 2001-2005 based on follow-up of patients through 2006. Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.


Leukemia

Washington State Residents by Gender
Incidence Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined and men and women separately, leukemia incidence rates remained level during the period from 1992 to 2007.

Washington State Residents by Gender
Death Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined, leukemia mortality rates decreased slightly by 0.8% per year during the period from 1992 to 2007.
- For men, leukemia mortality rates remained level during the period from 1992 to 2007.
- For women, leukemia mortality rates decreased steadily by 1.1% per year during the period from 1992 to 2007.

The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.


Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.

Leukemia

2005-2007 Washington State Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population




Age-adjusted rate per 100,000
 NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.
 Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2007 released 1/2010.

Leukemia

2005-2007 Washington State Residents

Average Annual Death Rates by County with rates in [] and 95% CI's []

Age-adjusted to 2000 US standard population



NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.

Liver

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using primary site ICD-03 code C22.0, excluding histology codes 9140, 9590-9989. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 codes 155.0, 155.2 (1992-1998), ICD-10 codes C22.0, C22.2-C22.4, C22.7, C22.9 (1999-2007).

2007 Washington State Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


| Incidence | | | |
|------------------------|------|--------|-------|
| | Male | Female | Total |
| All Reportable Cancers | 301 | 120 | 421 |
| 2007 Incidence Rates | 9.3 | 3.4 | 6.2 |
| 2006 National Rates | 9.8 | 3.2 | 6.3 |

| Death | | | |
|---------------------|------|--------|-------|
| | Male | Female | Total |
| Deaths | 210 | 73 | 283 |
| 2007 Death Rates | 6.7 | 2.0 | 4.2 |
| 2006 National Rates | 6.4 | 2.2 | 4.1 |

Stage at Diagnosis

2007 Washington State Residents

Total Incidence and Percents



Relative Survival Rates

2006 National 5-year rates¹

| | |
|----------|-----|
| In situ | n/a |
| Local | 27% |
| Regional | 9% |
| Distant | 3% |
| Unstaged | 6% |
| Overall | 14% |


n/a = not available

Figures include only invasive cancers

¹ 5-Year Relative Survival Rates for cases diagnosed 2001-2005 based on follow-up of patients through 2006. Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.


Liver

Washington State Residents by Gender
Incidence Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined, liver cancer incidence rates increased sharply by 5.2% per year during the period from 1992 to 2007.
- For men, liver cancer incidence rates increased sharply by 5.1% per year during the period from 1992 to 2007.
- For women, liver cancer incidence rates increased sharply by 4.5% during the period from 1992 to 2007.

Washington State Residents by Gender
Death Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined, liver cancer mortality rates increased steadily by 3.3% per year during the period from 1992 to 2007.
- For men, liver cancer mortality rates increased steadily by 3.6% per year during the period from 1992 to 2007.
- For women, liver cancer mortality rates increased steadily by 2.1% per year during the period from 1992 to 2007.

The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.


Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.

Liver

2005-2007 Washington State Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population




Age-adjusted rate per 100,000
 NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.
 Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2007 released 1/2010.

Liver

2005-2007 Washington State Residents

Average Annual Death Rates by County with rates in [] and 95% CI's [|]

Age-adjusted to 2000 US standard population



NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.

Lung and Bronchus

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using primary site ICD-03 codes C34.0-C34.9, excluding histology codes 9140, 9590-9989. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 codes 162.2-162.9 (1992-1998), ICD-10 code C34 (1999-2001).

2007 Washington State Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


| Incidence | | | |
|------------------------|-------|--------|-------|
| | Male | Female | Total |
| All Reportable Cancers | 1,915 | 1,939 | 3,856 |
| 2007 Incidence Rates | 68.0 | 56.2 | 61.0 |
| 2006 National Rates | 73.3 | 51.4 | 60.7 |

| Death | | | |
|---------------------|-------|--------|-------|
| | Male | Female | Total |
| Deaths | 1,600 | 1,568 | 3,168 |
| 2007 Death Rates | 57.7 | 45.0 | 50.4 |
| 2006 National Rates | 67.5 | 40.2 | 51.7 |

Stage at Diagnosis

2007 Washington State Residents

Total Incidence and Percents



Relative Survival Rates

2006 National 5-year rates¹

| | |
|----------|-----|
| In situ | 49% |
| Local | 50% |
| Regional | 22% |
| Distant | 3% |
| Unstaged | 8% |
| Overall | 16% |


n/c Number of cases <= 5

Figures include in situ and invasive cancers

¹ 5-Year Relative Survival Rates for cases diagnosed 2001-2005 based on follow-up of patients through 2006. Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.


Lung and Bronchus

Washington State Residents by Gender Incidence Rate Trends Age-adjusted to 2000 US standard population



- For both sexes combined, lung and bronchus cancer incidence rates remained level from 1992 to 1999; since then, however the rates have decreased steadily by 1.8% per year.
- For men, lung and bronchus cancer incidence rates decreased steadily by 1.4% per year from 1992 to 2005; since then, however the rates have remained level.
- For women, lung and bronchus cancer incidence rates increased steadily by 2.4% per year from 1992 to 1998; since then, however the rates have remained level.

Washington State Residents by Gender Death Rate Trends Age-adjusted to 2000 US standard population



- For both sexes combined, lung and bronchus cancer mortality rates decreased steadily by 1.3% per year during the period from 1992 to 2007.
- For men, lung and bronchus cancer mortality rates decreased steadily by 2.3% per year during the period from 1992 to 2007.
- For women, lung and bronchus cancer mortality rates remained level during the period from 1992 to 2007.

The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.


Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.

Lung and Bronchus

2005-2007 Washington State Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population



Age-adjusted rate per 100,000

NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.


Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2007 released 1/2010.

Lung and Bronchus

2005-2007 Washington State Residents

Average Annual Death Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population



NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.

Melanoma of the Skin

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using primary site ICD-03 codes C44.0-C44.9, including only histology codes 8720-8790. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 codes 172.0-172.9 (1992-1998), ICD-10 code C43 (1999-2007).

2007 Washington State Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


| Incidence | | | |
|------------------------|-------|--------|-------|
| | Male | Female | Total |
| All Reportable Cancers | 1,454 | 1,233 | 2,687 |
| 2007 Incidence Rates | 47.9 | 36.1 | 40.7 |
| 2006 National Rates | 42.1 | 26.6 | 33.0 |

| Death | | | |
|---------------------|------|--------|-------|
| | Male | Female | Total |
| Deaths | 139 | 67 | 206 |
| 2007 Death Rates | 4.6 | 1.9 | 3.1 |
| 2006 National Rates | 4.1 | 1.7 | 2.7 |

Stage at Diagnosis

2007 Washington State Residents

Total Incidence and Percents



Relative Survival Rates

2006 National 5-year rates¹


| | |
|----------|------|
| In situ | 100% |
| Local | 99% |
| Regional | 65% |
| Distant | 15% |
| Unstaged | 77% |
| Overall | 96% |

Figures include in situ and invasive cancers

¹ 5-Year Relative Survival Rates for cases diagnosed 2001-2005 based on follow-up of patients through 2006. Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.


Melanoma of the Skin

**Washington State Residents by Gender
Incidence Rate Trends**
Age-adjusted to 2000 US standard population



- For both sexes combined, melanoma incidence rates increased sharply by 11.3% per year from 1992 to 1995; rates increased steadily by 3.9% per year from 1995 to 2005; since 2005 the rates have remained level.
- For men, melanoma incidence rates increased sharply by 4.8% per year from 1992 to 2005; since then, however the rates have remained level.
- For women, melanoma incidence rates increased sharply by 12.5% per year from 1992 to 1995; rates increased steadily by 3.9% per year from 1995 to 2005; since 2005 the rates have remained level.

**Washington State Residents by Gender
Death Rate Trends**
Age-adjusted to 2000 US standard population



- For both sexes combined and men and women separately, melanoma mortality rates remained level during the period from 1992 to 2007.

The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.


Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.

Melanoma of the Skin

2005-2007 Washington State Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population




Age-adjusted rate per 100,000
 NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.
 Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2007 released 1/2010.

Melanoma of the Skin

2005-2007 Washington State Residents

Average Annual Death Rates by County with rates in [] and 95% CI's |—|

Age-adjusted to 2000 US standard population



NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.

Multiple Myeloma

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using all primary sites with histology codes 9731-9732, 9734. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 codes 203.0, 238.6 (1992-1998), ICD-10 codes C90.0, C90.2 (1999-2007).

2007 Washington State Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


| Incidence | | | |
|------------------------|------|--------|-------|
| | Male | Female | Total |
| All Reportable Cancers | 195 | 134 | 329 |
| 2007 Incidence Rates | 6.8 | 3.8 | 5.1 |
| 2006 National Rates | 6.6 | 4.3 | 5.3 |

| Death | | | |
|---------------------|------|--------|-------|
| | Male | Female | Total |
| Deaths | 124 | 88 | 212 |
| 2007 Death Rates | 4.7 | 2.5 | 3.5 |
| 2006 National Rates | 4.4 | 2.9 | 3.5 |

Stage at Diagnosis

2007 Washington State Residents

Total Incidence and Percents



Relative Survival Rates

2006 National 5-year rates¹

| | |
|----------|-----|
| In situ | n/a |
| Local | n/a |
| Regional | n/a |
| Distant | 39% |
| Unstaged | n/a |
| Overall | 39% |

n/a = not available


n/c Number of cases <= 5

Figures include only invasive cancers

¹ 5-Year Relative Survival Rates for cases diagnosed 2001-2005 based on follow-up of patients through 2006. Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.


Multiple Myeloma

Washington State Residents by Gender
Incidence Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined, multiple myeloma incidence rates remained level from 1992 to 1996; since then, however the rates have decreased steadily by 1.4% per year.
- For men, multiple myeloma incidence rates remained level from 1992 to 1996; since then, however the rates have decreased steadily by 1.4% per year.
- For women, multiple myeloma incidence rates decreased steadily by 1.0% per year during the period from 1992 to 2007.

Washington State Residents by Gender
Death Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined, multiple myeloma mortality rates remained level from 1992 to 2002; since then, however the rates have decreased sharply by 5.5% per year.
- For men, multiple myeloma mortality rates decreased steadily by 1.0% per year during the period from 1992 to 2007.
- For women, multiple myeloma mortality rates remained level during the period from 1992 to 2007.

The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.


Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.

Multiple Myeloma

2005-2007 Washington State Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population



Age-adjusted rate per 100,000


NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.
Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2007 released 1/2010.

Multiple Myeloma

2005-2007 Washington State Residents

Average Annual Death Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population



Age-adjusted rate per 100,000
 NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.

Non-Hodgkin Lymphoma

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry all sites with histology 9590-9596, 9670-9729; and except for sites C420, C421 and C424 all other sites with histology 9823 or 9827. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 codes 200.0-200.8, 202.0-202.2, 202.8-202.9 (1992-1998), ICD-10 codes C82-C85, C96.3 (1999-2007).

2007 Washington State Residents

Incidence and Death Summary by Sex


Counts and rates per 100,000 adjusted to the 2000 standard population

| Incidence | | | |
|------------------------|------|--------|-------|
| | Male | Female | Total |
| All Reportable Cancers | 755 | 599 | 1,355 |
| 2007 Incidence Rates | 25.4 | 17.1 | 20.8 |
| 2006 National Rates | 22.8 | 16.2 | 19.2 |

| Death | | | |
|---------------------|------|--------|-------|
| | Male | Female | Total |
| Deaths | 248 | 193 | 441 |
| 2007 Death Rates | 9.0 | 5.2 | 6.9 |
| 2006 National Rates | 8.4 | 5.4 | 6.7 |

Stage at Diagnosis

2007 Washington State Residents
Total Incidence and Percents



Relative Survival Rates

2006 National 5-year rates¹

| | |
|----------|-----|
| In situ | n/a |
| Local | n/a |
| Regional | n/a |
| Distant | n/a |
| Unstaged | n/a |
| Overall | 68% |


n/a = not available

Figures include only invasive cancers

¹ 5-Year Relative Survival Rates for cases diagnosed 2001-2005 based on follow-up of patients through 2006. Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.


Non-Hodgkin Lymphoma

Washington State Residents by Gender
Incidence Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined, incidence rates for Non-Hodgkin Lymphoma increased steadily by 1.5% per year from 1992 to 2004; since then, however the rates have remained level.
- For men, incidence rates for Non-Hodgkin Lymphoma increased sharply by 12.7% per year from 1992 to 1994; rates remained level from 1994 to 1998; rates increased steadily by 2.6% per year from 1998 to 2004; since 2004 the rates have remained level.
- For women, incidence rates for Non-Hodgkin Lymphoma increased steadily by 2.2% per year from 1992 to 2003; since then, however the rates have remained level.

Washington State Residents by Gender
Death Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined, mortality rates for Non-Hodgkin Lymphoma remained level from 1992 to 1999; since then, however the rates have decreased steadily by 3.7% per year.
- For men, mortality rates for Non-Hodgkin Lymphoma remained level from 1992 to 2002; since then, however the rates have decreased sharply by 6.5% per year.
- For women, mortality rates for Non-Hodgkin Lymphoma decreased steadily by 2.2% per year during the period from 1992 to 2007.

The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.


Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.

Non-Hodgkin Lymphoma

2005-2007 Washington State Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population




NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.
 Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2007 released 1/2010.

Non-Hodgkin Lymphoma

2005-2007 Washington State Residents

Average Annual Death Rates by County with rates in [] and 95% CI's |—|

Age-adjusted to 2000 US standard population



Age-adjusted rate per 100,000
 NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.

Oral Cavity and Pharynx

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using primary site ICD-03 codes C00.0-C14.8, excluding histology codes 9140, 9590-9989. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 codes 140.0-149.9 (1992-1998), ICD-10 codes C00-C14 (1999-2007).

2007 Washington State Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


| Incidence | | | |
|------------------------|------|--------|-------|
| | Male | Female | Total |
| All Reportable Cancers | 531 | 247 | 778 |
| 2007 Incidence Rates | 16.4 | 6.9 | 11.5 |
| 2006 National Rates | 15.3 | 6.2 | 10.4 |

| Death | | | |
|---------------------|------|--------|-------|
| | Male | Female | Total |
| Deaths | 125 | 69 | 194 |
| 2007 Death Rates | 4.2 | 1.9 | 3.0 |
| 2006 National Rates | 3.8 | 1.4 | 2.5 |

Stage at Diagnosis

2007 Washington State Residents

Total Incidence and Percents



Relative Survival Rates

2006 National 5-year rates¹


| | |
|----------|-----|
| In situ | 95% |
| Local | 84% |
| Regional | 55% |
| Distant | 30% |
| Unstaged | 53% |
| Overall | 62% |

Figures include in situ and invasive cancers

¹ 5-Year Relative Survival Rates for cases diagnosed 2001-2005 based on follow-up of patients through 2006. Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.


Oral Cavity and Pharynx

Washington State Residents by Gender
Incidence Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined, oral cavity and pharyngeal cancer incidence rates decreased slightly by 0.9% per year during the period from 1992 to 2007.
- For men, oral cavity and pharyngeal cancer incidence rates decreased slightly by 0.7% per year during the period from 1992 to 2007.
- For women, oral cavity and pharyngeal cancer incidence rates decreased steadily by 1.5% per year during the period from 1992 to 2007.

Washington State Residents by Gender
Death Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined and men and women separately, oral cavity and pharynx cancer mortality rates remained level during the period from 1992 to 2007.

The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.


Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.

Oral Cavity and Pharynx

2005-2007 Washington State Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population




NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.
 Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2007 released 1/2010.

Oral Cavity and Pharynx

2005-2007 Washington State Residents

Average Annual Death Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population



Age-adjusted rate per 100,000
 NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.

Ovary

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using primary site ICD-03 code C56.9, excluding histology codes 9140, 9590-9989. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 code 183.0 (1992-1998), ICD-10 code C56 (1999-2007).

2007 Washington State Female Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


| Incidence | | | |
|------------------------|------|--------|-------|
| | Male | Female | Total |
| All Reportable Cancers | n/a | 459 | n/a |
| 2007 Incidence Rates | n/a | 12.7 | n/a |
| 2006 National Rates | n/a | 12.7 | n/a |

| Death | | | |
|---------------------|------|--------|-------|
| | Male | Female | Total |
| Deaths | n/a | 371 | n/a |
| 2007 Death Rates | n/a | 10.3 | n/a |
| 2006 National Rates | n/a | 8.5 | n/a |

Stage at Diagnosis

2007 Washington State Female Residents

Total Incidence and Percents



Relative Survival Rates

2006 National 5-year rates¹

| | |
|----------|-----|
| In situ | 97% |
| Local | 93% |
| Regional | 71% |
| Distant | 31% |
| Unstaged | 25% |
| Overall | 46% |

n/c Number of cases <= 5

Figures include in situ and invasive cancers


¹ 5-Year Relative Survival Rates for cases diagnosed 2001-2005 based on follow-up of patients through 2006.

n/a Not available for sex-specific sites.

Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.


Ovary

Washington State Female Residents
Incidence Rate Trends
 Age-adjusted to 2000 US standard population



- For women, ovarian cancer incidence rates remained level from 1992 to 2002; since then, however the rates have decreased sharply by 4.9% per year.

Washington State Female Residents
Death Rate Trends
 Age-adjusted to 2000 US standard population



- For women, ovarian cancer mortality rates remained level during the period from 1992 to 2007.

The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.


Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry- Incidence data for diagnosis years 1992 to 2007 released in January 2010.

Ovary

2005-2007 Washington State Female Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population




Age-adjusted rate per 100,000
 NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.
 Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2007 released 1/2010.

Ovary

2005-2007 Washington State Female Residents

Average Annual Death Rates by County with rates in [] and 95% CI's

Age-adjusted to 2000 US standard population



Age-adjusted rate per 100,000
 NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.

Pancreas

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using primary site ICD-0 codes C25.0-C25.9, excluding histology codes 9140, 9590-9989. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 codes 157.0-157.9 (1992-1998), ICD-10 code C25 (1999-2007).

2007 Washington State Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


| Incidence | | | |
|------------------------|------|--------|-------|
| | Male | Female | Total |
| All Reportable Cancers | 384 | 388 | 772 |
| 2007 Incidence Rates | 13.4 | 10.7 | 11.9 |
| 2006 National Rates | 13.2 | 10.5 | 11.7 |

| Death | | | |
|---------------------|------|--------|-------|
| | Male | Female | Total |
| Deaths | 363 | 343 | 706 |
| 2007 Death Rates | 12.7 | 9.4 | 10.9 |
| 2006 National Rates | 12.4 | 9.5 | 10.8 |

Stage at Diagnosis

2007 Washington State Residents

Total Incidence and Percents



Relative Survival Rates

2006 National 5-year rates¹

| | |
|----------|-----|
| In situ | 84% |
| Local | 22% |
| Regional | 9% |
| Distant | 2% |
| Unstaged | 4% |
| Overall | 6% |


n/c Number cases <= 5

Figures include in situ and invasive cancers

¹ 5-Year Relative Survival Rates for cases diagnosed 2001-2005 based on follow-up of patients through 2006. Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.


Pancreas

Washington State Residents by Gender
Incidence Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined, pancreatic cancer incidence rates increased slightly by 0.7% per year during the period from 1992 to 2007.
- For men, pancreatic cancer incidence rates increased slightly by 0.9% per year during the period from 1992 to 2007.
- For women, pancreatic cancer incidence rates increased steadily by 0.5% per year during the period from 1992 to 2007.

Washington State Residents by Gender
Death Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined and men and women separately, pancreatic cancer mortality rates remained level during the period from 1992 to 2007.

The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.


Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.

Pancreas

2005-2007 Washington State Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population




Age-adjusted rate per 100,000
 NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.
 Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2007 released 1/2010.

Pancreas

2005-2007 Washington State Residents

Average Annual Death Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population



Age-adjusted rate per 100,000
 NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.

Prostate

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using primary site ICD-03 code C61.9, excluding histology codes 9140, 9590-9989. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 code 185 (1992-1998), ICD-10 code C61 (1999-2007).

2007 Washington State Male Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


| Incidence | | | |
|------------------------|-------|--------|-------|
| | Male | Female | Total |
| All Reportable Cancers | 4,747 | n/a | n/a |
| 2007 Incidence Rates | 158.4 | n/a | n/a |
| 2006 National Rates | 154.0 | n/a | n/a |

| Death | | | |
|---------------------|------|--------|-------|
| | Male | Female | Total |
| Deaths | 635 | n/a | n/a |
| 2007 Death Rates | 25.2 | n/a | n/a |
| 2006 National Rates | 23.6 | n/a | n/a |

Stage at Diagnosis

2007 Washington State Male Residents

Total Incidence and Percents



Relative Survival Rates

2006 National 5-year rates¹

| | |
|----------|------|
| In situ | 96% |
| Local | 100% |
| Regional | 100% |
| Distant | 30% |
| Unstaged | 74% |
| Overall | 100% |

n/c Number of cases <= 5

Figures include in situ and invasive cancers


¹ 5-Year Relative Survival Rates for cases diagnosed 2001-2005 based on follow-up of patients through 2006.

n/a Not available for sex-specific sites.

Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.


Prostate

**Washington State Male Residents
Incidence Rate Trends**
Age-adjusted to 2000 US standard population



- For men, prostate cancer incidence rates decreased sharply by 12.6% per year from 1992 to 1995, rates increased sharply by 4.8% per year from 1995 to 1999; since 1999 the rates have decreased steadily by 2.0% per year.

**Washington State Male Residents
Death Rate Trends**
Age-adjusted to 2000 US standard population



- For men, prostate cancer mortality rates decreased sharply by 4.9% per year from 1992 to 1998; since 1998 the rates have decreased steadily by 2.0% per year.

The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.


Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.

Prostate

2005-2007 Washington State Male Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population




Age-adjusted rate per 100,000
 NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.
 Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2007 released 1/2010.

Prostate

2005-2007 Washington State Male Residents

Average Annual Death Rates by County with rates in [] and 95% CI 's |

Age-adjusted to 2000 US standard population



Age-adjusted rate per 100,000

NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.

Soft Tissue including Heart

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using primary site ICD-03 codes C38.0, C47.0-C47.9, C49.0-C49.9, excluding histology codes 9140, 9590-9989. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 codes 164.1, 171.0-171.9 (1992-1998), ICD-10 codes C47, C49, C38.0, C45.2 (1999-2007).

2007 Washington State Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


| Incidence | | | |
|------------------------|------|--------|-------|
| | Male | Female | Total |
| All Reportable Cancers | 118 | 93 | 211 |
| 2007 Incidence Rates | 3.9 | 2.6 | 3.2 |
| 2006 National Rates | 3.7 | 2.5 | 3.0 |

| Death | | | |
|---------------------|------|--------|-------|
| | Male | Female | Total |
| Deaths | 35 | 36 | 71 |
| 2007 Death Rates | 1.3 | 1.0 | 1.1 |
| 2006 National Rates | 1.5 | 1.1 | 1.3 |

Stage at Diagnosis

2007 Washington State Residents

Total Incidence and Percents



Relative Survival Rates

2006 National 5-year rates¹

| | |
|----------|-----|
| In situ | n/a |
| Local | 85% |
| Regional | 63% |
| Distant | 17% |
| Unstaged | 55% |
| Overall | 67% |

n/a = not available


n/c Number of cases <= 5

Figures include only invasive cancers

¹ 5-Year Relative Survival Rates for cases diagnosed 2001-2005 based on follow-up of patients through 2006. Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.


Soft Tissue including Heart

Washington State Residents by Gender
Incidence Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined and men and women separately, incidence rates for soft tissue including heart remained level during the period from 1992 to 2007.

Washington State Residents by Gender
Death Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined, mortality rates for soft tissue including heart decreased steadily by 3.0% per year during the period from 1992 to 2007.
- For men, mortality rates for soft tissue including heart decreased steadily by 2.5% per year during the period from 1992 to 2007.
- For women, mortality rates for soft tissue including heart decreased steadily by 3.4% per year during the period from 1992 to 2007.

The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.


Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.

Soft Tissue including Heart

2005-2007 Washington State Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's

Age-adjusted to 2000 US standard population




Age-adjusted rate per 100,000
 NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.
 Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2007 released 1/2010.

Soft Tissue including Heart

2005-2007 Washington State Residents

Average Annual Death Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population



NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.

Stomach

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using primary site ICD-03 codes C16.0-C16.9, excluding histology codes 9140, 9590-9989. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 codes 151.0-151.9 (1992-1998), ICD-10 code C16 (1999-2007).

2007 Washington State Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


| Incidence | | | |
|------------------------|------|--------|-------|
| | Male | Female | Total |
| All Reportable Cancers | 252 | 146 | 398 |
| 2007 Incidence Rates | 8.5 | 4.1 | 6.1 |
| 2006 National Rates | 10.5 | 5.4 | 7.6 |

| Death | | | |
|---------------------|------|--------|-------|
| | Male | Female | Total |
| Deaths | 121 | 93 | 214 |
| 2007 Death Rates | 4.3 | 2.7 | 3.3 |
| 2006 National Rates | 5.0 | 2.6 | 3.7 |

Stage at Diagnosis

2007 Washington State Residents

Total Incidence and Percents



Relative Survival Rates

2006 National 5-year rates¹


| | |
|----------|-----|
| In situ | 72% |
| Local | 61% |
| Regional | 28% |
| Distant | 4% |
| Unstaged | 18% |
| Overall | 27% |

Figures include in situ and invasive cancers

¹ 5-Year Relative Survival Rates for cases diagnosed 2001-2005 based on follow-up of patients through 2006. Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.


Stomach

Washington State Residents by Gender
Incidence Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined, stomach cancer incidence rates decreased steadily by 1.1% per year during the period from 1992 to 2007.
- For men, stomach cancer incidence rates decreased steadily by 1.6% per year during the period from 1992 to 2007.
- For women, stomach cancer incidence rates remained level during the period from 1992 to 2007.

Washington State Residents by Gender
Death Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined, stomach cancer mortality rates decreased steadily by 3.0% per year during the period from 1992 to 2007.
- For men, stomach cancer mortality rates decreased steadily by 3.9% per year during the period from 1992 to 2007.
- For women, stomach cancer mortality rates decreased steadily by 1.8% per year during the period from 1992 to 2007.

The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.


Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.

Stomach

2005-2007 Washington State Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population




Age-adjusted rate per 100,000
 NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.
 Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2007 released 1/2010.

Stomach

2005-2007 Washington State Residents

Average Annual Death Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population



Age-adjusted rate per 100,000

NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.

Testis

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using primary site ICD-03 codes C62.0-C62.9, excluding histology codes 9140, 9590-9989. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 codes 186.0-186.9 (1992-1998), ICD-10 code C62 (1999-2007).

2007 Washington State Male Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


| Incidence | | | |
|------------------------|------|--------|-------|
| | Male | Female | Total |
| All Reportable Cancers | 194 | n/a | n/a |
| 2007 Incidence Rates | 5.9 | n/a | n/a |
| 2006 National Rates | 5.3 | n/a | n/a |

| Death | | | |
|---------------------|------|--------|-------|
| | Male | Female | Total |
| Deaths | 9 | n/a | n/a |
| 2007 Death Rates | 0.3 | n/a | n/a |
| 2006 National Rates | 0.2 | n/a | n/a |

Stage at Diagnosis

2007 Washington State Male Residents

Total Incidence and Percents



Relative Survival Rates

2006 National 5-year rates¹

| | |
|----------|------|
| In situ | 100% |
| Local | 99% |
| Regional | 96% |
| Distant | 71% |
| Unstaged | 84% |
| Overall | 95% |

n/c Number of cases <= 5

Figures include only invasive cancers


¹ 5-Year Relative Survival Rates for cases diagnosed 2001-2005 based on follow-up of patients through 2006.

n/a Not available for sex-specific sites.

Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.


Testis

Washington State Male Residents
Incidence Rate Trends
 Age-adjusted to 2000 US standard population



- For men, testicular cancer incidence rates remained level during the period from 1992 to 2007.

Washington State Male Residents
Death Rate Trends
 Age-adjusted to 2000 US standard population



- For men, testicular cancer mortality rates remained level during the period from 1992 to 2007.

The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.


Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.

Testis

2005-2007 Washington State Male Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population




Age-adjusted rate per 100,000
 NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.
 Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2007 released 1/2010.

Testis

2005-2007 Washington State Male Residents

Average Annual Death Rates by County with rates in [] and 95% CI 's |

Age-adjusted to 2000 US standard population



Age-adjusted rate per 100,000
 NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases ≤ 5.

Thyroid

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using primary site ICD-03 code C73.9, excluding histology codes 9140, 9590-9989. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 code 193 (1992-1998), ICD-10 code C73 (1999-2007).

2007 Washington State Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


| Incidence | | | |
|------------------------|------|--------|-------|
| | Male | Female | Total |
| All Reportable Cancers | 188 | 540 | 729 |
| 2007 Incidence Rates | 5.8 | 16.2 | 11.0 |
| 2006 National Rates | 5.4 | 15.7 | 10.6 |

| Death | | | |
|---------------------|------|--------|-------|
| | Male | Female | Total |
| Deaths | 25 | 23 | 48 |
| 2007 Death Rates | 0.9 | 0.6 | 0.8 |
| 2006 National Rates | 0.5 | 0.5 | 0.5 |

Stage at Diagnosis

2007 Washington State Residents

Total Incidence and Percents



Relative Survival Rates

2006 National 5-year rates¹

| | |
|----------|------|
| In situ | n/a |
| Local | 100% |
| Regional | 97% |
| Distant | 59% |
| Unstaged | 89% |
| Overall | 97% |


n/a = not available

Figures include in situ and invasive cancers

¹ 5-Year Relative Survival Rates for cases diagnosed 2001-2005 based on follow-up of patients through 2006. Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.


Thyroid

Washington State Residents by Gender
Incidence Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined, thyroid cancer incidence rates increased sharply by 5.2% per year during the period from 1992 to 2007.
- For men, thyroid cancer incidence rates increased sharply by 4.9% per year during the period from 1992 to 2007.
- For women, thyroid cancer incidence rates increased sharply by 5.3% per year during the period from 1992 to 2007.

Washington State Residents by Gender
Death Rate Trends
 Age-adjusted to 2000 US standard population



- For both sexes combined, men and women separately, thyroid cancer mortality rates remained level during the period from 1992 to 2007.

The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.


Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.

Thyroid

2005-2007 Washington State Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population




Age-adjusted rate per 100,000
 NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.
 Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2007 released 1/2010.

Thyroid

2005-2007 Washington State Residents

Average Annual Death Rates by County with rates in [] and 95% CI's []

Age-adjusted to 2000 US standard population



Age-adjusted rate per 100,000
 NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.

Vulva (Female)

DATA DEFINITIONS: Incidence data were obtained from the Washington State Cancer Registry using primary site ICD-03 code C51.0-C51.9, excluding histology codes 9140, 9590-9989. Death data were obtained from Washington State death certificates using the underlying cause of death ICD-9 code 184.1-184.4 (1992-1998), ICD-10 code C51 (1999-2007).

2007 Washington State Female Residents

Incidence and Death Summary by Sex

Counts and rates per 100,000 adjusted to the 2000 standard population


| Incidence | | | |
|------------------------|------|--------|-------|
| | Male | Female | Total |
| All Reportable Cancers | n/a | 233 | n/a |
| 2007 Incidence Rates | n/a | 6.7 | n/a |
| 2006 National Rates | n/a | 6.1 | n/a |

| Death | | | |
|---------------------|------|--------|-------|
| | Male | Female | Total |
| Deaths | n/a | 11 | n/a |
| 2007 Death Rates | n/a | 0.3 | n/a |
| 2006 National Rates | n/a | 0.5 | n/a |

Stage at Diagnosis

2007 Washington State Female Residents

Total Incidence and Percents



Relative Survival Rates

2006 National 5-year rates¹

| | |
|----------|-----|
| In situ | 98% |
| Local | 90% |
| Regional | 56% |
| Distant | 11% |
| Unstaged | 60% |
| Overall | 90% |

n/c Number of cases <= 5

Figures include in situ and invasive cancers


¹ 5-Year Relative Survival Rates for cases diagnosed 2001-2005 based on follow-up of patients through 2006.

n/a Not available for sex-specific sites.

Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry-Incidence data for diagnosis years 1992 to 2007 released in January 2010.


Vulva (Female)

Washington State Female Residents
Incidence Rate Trends
 Age-adjusted to 2000 US standard population



- For women, vulva cancer incidence rates remained level during the period from 1992 to 2007.

Washington State Female Residents
Death Rate Trends
 Age-adjusted to 2000 US standard population



- For women, vulva cancer mortality rates remained level during the period from 1992 to 2007.

The coding for causes of death changed in 1999. These changes do not substantively affect the trends for cancer deaths.


Washington State Cancer Incidence Data: Washington State Department of Health, Washington State Cancer Registry- Incidence data for diagnosis years 1992 to 2007 released in January 2010.

Vulva (Female)

2005-2007 Washington State Female Residents

Average Annual Incidence Rates by County with rates in [] and 95% CI's |

Age-adjusted to 2000 US standard population




Age-adjusted rate per 100,000
 NOTE: Rates and confidence intervals are not computed when the 2005-2007 total number of new cases <= 5.
 Washington State Cancer Registry Database: Incidence data for diagnosis year 1992 to 2007 released 1/201

Vulva (Female)

2005-2007 Washington State Female Residents

Average Annual Death Rates by County with rates in [] and 95% CI's

Age-adjusted to 2000 US standard population



Age-adjusted rate per 100,000
 NOTE: Rates and confidence intervals are not computed when the
 2005-2007 total number of new cases ≤ 5