

**WASHINGTON DEPARTMENT OF ECOLOGY  
 ENVIRONMENTAL ASSESSMENT PROGRAM  
 FRESHWATER MONITORING UNIT  
 STREAM DISCHARGE TECHNICAL NOTES  
 MANUAL STAGE HEIGHT STATION**

**STATION ID:** 45B070  
**STATION NAME:** Icicle Creek near Leavenworth  
**WATER YEAR:** 2010  
**AUTHOR:** Tyler W. Burks

**Introduction**

Watershed Description

Icicle Creek originates in the snowfields of the eastern slopes of the central Cascade Mountain range and flows into the Wenatchee River at the City of Leavenworth. The watershed is bounded by both the Stuart Range and the Chiwaukum Mountains. Land cover above the gage consists of predominantly coniferous forest but also includes alpine shrubland, montane grassland, bedrock/talus slopes, and riparian woodlands of the Wenatchee National Forest and Alpine Lakes Wilderness Area. Mean annual precipitation across the watershed above this gage location is 82.3 inches (U.S. Weather Bureau, 1965).

Gage Location

Observation of the manual stage height stream gaging station on Icicle Creek began on May 3, 2007. The gage is located at the East Leavenworth Road bridge on the right bank, approximately one half mile downstream of the Leavenworth National Fish Hatchery Complex.

Table 1.

Drainage Area (square miles)	211 (USGS, 2012)
Latitude (degrees, minutes, seconds)	47° 33' 49" N
Longitude (degrees, minutes, seconds)	120° 40' 04" W
Primary Gage Index Type	Staff Gage
Secondary Gage Index Type	Tape Down

## Error Analysis

Overall Rating Error Percentage	10.1%
---------------------------------	-------

## Rating Table(s)

Table 2. Rating Table Summary

Rating Table No.	1		
Period of Ratings	5/2007-10/2012		
Range of Ratings (cfs)	47-9230		
No. of Defining Measurements	38		
Rating Error (%)	10.1%		

## Narrative

Rating Table #1 applies to the current period of record at this station. It has evolved slightly with the addition of new measurements, but, overall has continued to be persistent. The rating has been logarithmically extended to cover one half the lowest measured discharge and two times the highest measured discharge.

## Discrete Flow Record

Table 3. Discrete Flow Record Summary

Number of Discrete Stage Readings	38	
Maximum Observed Stage (feet) and Date	7.59	06/09/2010
Maximum Predicted Discharge (cfs) and Date	2890	06/09/2010
Minimum Observed Stage (feet) and Date	1.88	10/08/2009
Minimum Predicted Discharge (cfs) and Date	78	10/08/2009
Range of Stage (feet) and Discharge (cfs)	1.88 to 7.59	78 to 2890

## Narrative

Funding for the Chelan County Natural Resources Department to make staff gage observations was eliminated prior to water year 2010. The data set was augmented by USFWS volunteers Dick Rieman and Fred Wurster, who collected observations during late winter and early spring. Seven discharge measurements were made during the water year, ranging from 205 to 2890 cfs. Based on continuous data collected by the USGS at Icicle Creek above Snow Creek, near Leavenworth (12458000), the peak occurred on June 3, 2010. Additionally, the minimum discharge for the water year occurred on October 11-13, 2009.

## Modeled Discharge

Table 4. Model Summary

Model Type (Slope conveyance, other, none)	---
Range of Modeled Stage (feet)	---
Range of Modeled Discharge (cfs)	---
Valid Period for Model	---
Model Confidence	---

## Surveys

Table 5. Survey Type and Date (station, cross section, longitudinal)

Type	Date
Station Levels	10/22/2008

## Activities Completed

--